

[JJ] Evening Poster | M (Multidisciplinary and Interdisciplinary) | M-IS Intersection

[M-IS10]Paleoclimatology and paleoceanography

convener:Yusuke Okazaki(Department of Earth and Planetary Sciences, Graduate School of Science, Kyushu University), Atsuhiko Isobe(Research Institute for Applied Mechanics, Kyushu University), Akihisa Kitamura(静岡大学理学部地球科学教室, 共同), Masaki Sano(Faculty of Human Sciences, Waseda University)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Past environmental changes and events at multi-decadal to tectonic timescale toward an understanding of Earth climate system by an integration of terrestrial and marine proxy studies and numerical modeling will be discussed. We welcome a variety of paleo-environmental studies from a wide range of background. In particular, a series of presentations relating to the Anthropocene will be planned. This is a merged session of A-OS31 "Linkage between oceanography and paleoceanography in marginal, shelf and coastal oceans" and M-IS23 "Paleoclimatology and paleoceanography" sessions at JPGU 2017. We hope that this session will provide an opportunity to promote communication between participants from multidisciplinary field.

[MIS10-P15]Abrupt warming at the onset of last deglaciation inferred from isotopic composition of fluid inclusions from a stalagmite in Okinawa

*Koki Nakamura¹, Satoru Mishima¹, Kanako Omine¹, Ryuji Asami², Jin-Ping Chen³, Chuan-Chou Shen³, Ryu Uemura¹ (1.University of the Ryukyus, 2.Tohoku University, 3.Dept. of Geosci., National Taiwan University)

Keywords:stalagmite, fluid inclusions, stable isotope, last deglaciation

Millennial scale abrupt climate changes have been occurred during the last deglaciation. Stalagmites is one of the most important archive to reconstruct terrestrial climate change in low- and mid-latitudes areas because of their high-accuracy of U-Th dating. However, interpretation of oxygen isotopic ratio ($\delta^{18}\text{O}_{\text{ca}}$) of calcite is not straightforward because it is controlled by two factors; $\delta^{18}\text{O}$ of dripwater and temperature at calcite formation.

Oxygen isotope ratio ($\delta^{18}\text{O}_{\text{fi}}$) and hydrogen isotopic ratio ($\delta\text{D}_{\text{fi}}$) of fluid-inclusions water preserve past isotopic ratios of drip water. Thus, they are useful to reconstruct not only the isotope composition of past rain water but also changes in temperature. In this study, we analyzed the $\delta^{18}\text{O}_{\text{fi}}$ and $\delta\text{D}_{\text{fi}}$ in the fluid inclusions and the $\delta^{18}\text{O}_{\text{ca}}$ of a stalagmite. A stalagmite, HSN2, was collected in Hoshino cave in Minami Daito Island, Okinawa, Japan. The isotope compositions of fluid inclusion water were measured using a semi-automated version of our fluid inclusion analysis system (Uemura et al., 2016). The $\delta^{18}\text{O}_{\text{ca}}$ of calcite was measured using Gas-bench CF-IRMS (Delta V advantage). U-Th dates was measured at National Taiwan University. The stalagmite HSN 2 has grown continuously from about 21,800 to about 13,400 years before present, and thus covering a important transition period between the Last Glacial Maximum (LGM), Heinrich stadial 1 (H1), and Bølling-Allerød (BA). The temperature variation estimated from $\delta^{18}\text{O}_{\text{fi}}$ and $\delta^{18}\text{O}_{\text{ca}}$ shows significant increase in temperature from H1 to BA.