

---

[JJ] Evening Poster | M (Multidisciplinary and Interdisciplinary) | M-IS Intersection

## [M-IS11]tsunami deposit

convener:Tetsuya Shinozaki(Center for Research in Isotopes and Environmental Dynamics (CRIED), University of Tsukuba), Takashi Chiba(Maritime Disaster Prevention Center), Daisuke Ishimura(首都大学東京大学院都市環境科学研究科地理学教室)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

The 2011 off the Pacific coast of Tohoku Earthquake and tsunami have an influence on the development of tsunami deposit research. After the tsunami, a lot of findings have been reported on various research fields. However, identification criteria of the tsunami deposit are not yet established. Moreover, it is still uncertain how to use the tsunami deposit in the risk assessment. In this session, we welcome researches from all aspect of sedimentary records of modern and paleo tsunamis both onshore and offshore, and numerical and experimental modeling studies for risk assessment. In addition, we also welcome other event deposits, such as flooding and storm surge, that they are considered to be important for discrimination of tsunami deposit.

---

## [MIS11-P09]Geochemical detection of the historical tsunami deposit

\*Yusuke Ogata<sup>1</sup>, Kazuhisa Goto<sup>2</sup>, Tetsuya Shinozaki<sup>3</sup>, Minoru Ikehara<sup>4</sup>, Catherine Chagué<sup>5</sup>, Takao Kawamata<sup>6</sup>  
(1.Department of Earth Sciences, Graduate School of Science, Tohoku University, 2.International Research Institute for Disaster Science (IRIDeS), Tohoku University, 3.Center for Research in Isotopes and Environmental Dynamics (CRIED), University of Tsukuba, 4.Center for Advanced Marine Core Research, Kochi University, 5.University of New South Wales (UNSW), 6.Educational Committee, City of Iwanuma)

Keywords:tsunami deposits, geochemistry

Historical records reveal that the Sendai and Ishinomaki plains were devastated by the AD869 Jogan and AD1611 Keicho tsunamis, and that the AD1454 Kyotoku tsunami might have also affected these plains. Radiocarbon dating is not straightforward for estimating ages in the medieval period, and thus the discrimination between the tsunami deposits left by the AD1611 and AD1454 events is difficult. Another issue is that the evidence of minor tsunamis might be hard to decipher in the geological record. Geochemistry is a useful tool which can help solve these issues, and its value has been increasingly recognized in paleo-tsunami and paleo-environmental research. Multi-proxy geochemical analyses in particular enable a more accurate interpretation of the geological record. In this study, we analyzed a core excavated from Iwanuma City, Miyagi Prefecture, using non-destructive X-ray fluorescence (ITRAX), as well as TC, TOC, TN and C/N ratio to identify the possible tsunami deposits and to reconstruct the paleo-environmental changes during the last thousand years, in combination with information from historical documents and diatom assemblages. We detected geochemical signatures that suggest a seawater inundation event, possibly a tsunami or storm, beneath the possible AD1611 tsunami deposit.