[JJ] Evening Poster | M (Multidisciplinary and Interdisciplinary) | M-IS Intersection

[M-IS11]tsunami deposit

convener:Tetsuya Shinozaki(Center for Research in Isotopes and Environmental Dynamics (CRiED), University of Tsukuba), Takashi Chiba(Maritime Disaster Prevention Center), Daisuke Ishimura(首都大学東 京大学院都市環境科学研究科地理学教室)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe) The 2011 off the Pacific coast of Tohoku Earthquake and tsunami have an influence on the development of tsunami deposit research. After the tsunami, a lot of findings have been reported on various research fields. However, identification criteria of the tsunami deposit are not yet established. Moreover, it is still uncertain how to use the tsunami deposit in the risk assessment. In this session, we welcome researches from all aspect of sedimentary records of modern and paleo tsunamis both onshore and offshore, and numerical and experimental modeling studies for risk assessment. In addition, we also welcome other event deposits, such as flooding and storm surge, that they are considered to be important for discrimination of tsunami deposit.

[MIS11-P15]Geological evidence for extreme wave events on the Sagara coastal lowland facing the Tōkai segment of the Nankai-Suruga Trough

Philipp Kempf¹, Osamu Fujiwara³, Yosuke Miyairi⁴, Yusuke Yokoyama⁴, Atsunori Nakamura³, Ed Garrett^{2,1}, Marc De Batist⁵, *Vanessa Mary An Heyvaert^{1,5}, QuakeRecNankai team (1.Geological Survey of Belgium, OD Earth and History of Life, Royal Belgian Institute of Natural Sciences, Jennerstraat 13, 1000 Brussels, Belgium, 2.Department of Geography and Institute of Hazard, Risk and Resilience, Durham University, United Kingdom, 3.Geological Survey of Japan, National Institute of Advanced Industrial Science and Technology, Japan, 4. Atmosphere and Ocean Research Institute, University of Tokyo, Japan, 5.Department of Geology and Soil Science, Ghent University, Belgium) Keywords:tsunami history, floodplain sediments, XRF scanning, CT-scans

Located close to Japan's densest concentrations of people and industry, the easternmost region of the Nankai-Suruga subduction zone has long been the focus of attempts to forecast future earthquakes. Historical records of the region attest to the occurrence of great earthquakes and subsequent tsunamis that may have originated from the Tōkai segment. However, past rupture zone extents and recurrence intervals remain poorly understood. Coastal stratigraphy has the potential to record the occurrence of both tsunami inundation and coseismic vertical land-level change over timescales far exceeding the historical record, with important implications for refining understanding of future hazards. Here we present results from an extensive coring survey of the lower reaches of the floodplain of the Sagara River, close to the town of Sagara, Shizuoka Prefecture. The site lies at an altitude of ~1 – 5 m and is within the anticipated inundation zone of future worst-case tsunami scenarios. Typhoon-driven storm surges and river floods are also likely to have inundated the site, complicating the interpretation of potential tsunami deposits. Using X-ray fluorescence scanning, X-ray computer tomography scans, multi-sensor core logs and radiocarbon dates, we evaluate sedimentary processes to be able to distinguish between extreme wave events and fluvial deposits. Where possible, we assess methods to differentiate between storm surges and tsunami deposits. Finally, we evaluate the potential for the site to provide a long and continuous record of extreme wave events and highlight the probable influence of changing thresholds of evidence creation and preservation over time.