[JJ] Evening Poster | M (Multidisciplinary and Interdisciplinary) | M-IS Intersection

[M-IS17]Gas hydrates in environmental-resource sciences

convener:Hitoshi Tomaru(Department of Earth Sciences, Chiba University), Akihiro Hachikubo(Kitami Institute of Technology), Atsushi Tani(神戸大学 大学院人間発達環境学研究科, 共同), Shusaku Goto(Institute for Geo-Resources and Environment National Institute of Advanced Industrial Science and Technology)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe) An increasing number of researches focusing on natural gas hydrates has recently been conducted from the environmental, material, and resource scientific viewpoints. This session aims to share and discuss the latest research results to understand and examine the nature and potential of gas hydrates in the past-present-future of the Earth. Because the researches on gas hydrates are interdisciplinary, broad topics from field and experimental researches, modeling, etc. will be presented in this session.

[MIS17-P10]Isotopic gas and microbial analyses of gas hydratebearing core in subsea floor sediments,offshore Sakhalin

*Kazui Sukawa¹, Akihiro Hachikubo¹, Hirotoshi Sakagami¹, Hirotsugu Minami¹, Satoshi Yamashita¹, Nobuo Takahashi¹, Hitoshi Shoji¹, Young K Jin², Boris Baranov³, Anatoly Obzhirov⁴, Masaaki Konishi¹ (1. Kitami Institute of Technology, 2.Korea Polar Research Institute, Korea, 3.P. P. Shirshov Institute of Oceanology, RAS, Russia, 4. V. I. Il'ichev Pacific Oceanological Institute FEB RAS, Russia)

Gas hydrates (GH) are crystalline water-based solids including hydrocarbons under high pressure and low temperature. Methane as the major gases of GH are considered to generate from organic matters buried under the benthic sediments by theological clacking reactions and / or by methanogenesis. Isotope compositions of methane are considered to record the gas-generating manner, because microbes selectively metabolitze ¹²C, in the field of geochemistry. In addition, microbial compositions of the GH sediments have been often studied all over the world. However, combination study by both isotopic gas and microbial diversities analyses is insufficient to understand the gas-generating behavior. In this study, using microbial and isotopic analyses, we examined a unique GH-burying core; obtained in the western Sakhalin slope off Sakhalin Island during Japan-Russia-Korea joint projects (SSGH projects).

The GH-gas confirmed as thermogenic gas on the basis of Barnard plots, and including higher H₂S and ethane than the other cores using principal component analysis of gas compositions. According to DNA-based microbial composition analysis, the Archaea abundance reduced along with the sediment depth. Under 20 cmbsf, MBG-B, MBG-D and Woesearchaeota (DHVEG-6) were dominated in Archaea. No methanogen was detected. Heterotrophic Chloroflexi and Candidate Atribacteria were major bacteria at a depth of 100 to 260 cm. It had been reported that Atribacteria decompose organic matters and generate ethanol and acids, by single cell genomics and metagenomics strategies. Since the low-molecules can be metabolized by methanogens, the Atribacteria-dominated bacteria composition may exhibit the possibility of methanogens, but methanogen was not detected. Low resolution of cloning-based DNA analysis might interrupt to detect methanogen. We, therefore, attempt to perform amplicon sequencing by Illumina the next generation sequencer.