
[JJ] Evening Poster | M (Multidisciplinary and Interdisciplinary) | M-IS Intersection

[M-IS18] Aqua planetology

convener: Yasuhito Sekine (Department of Earth and Planetary Science, University of Tokyo), Tomohiro Usui (Earth-Life Science Institute, Tokyo Institute of Technology), Hidenori Genda (東京工業大学 地球生命研究所, 共同), Takazo Shibuya (Japan Agency for Marine-Earth Science and Technology)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

This proposed session covers a wide range of topics related to aqua planetology, including chemical reactions involving water on planetary bodies, water-rock reactions within planetesimals, distribution of water in the early Solar system and the origin of water on terrestrial planets, hydrological and biogeochemical processes on Earth, geochemical cycles and habitability on Mars and icy moons, exploration of water in the solar system, and theory to understand how to build a habitable aqua planet.

[MIS18-P10] A probabilistic approach to deriving Ceres average surface composition from Dawn VIR data

*Hiroyuki Kurokawa¹, Bethany L Ehlmann^{2,3}, Eleonora Ammannito⁴, Maria Cristina De Sanctis⁵, Mathieu G. A. Lapotre⁶, Tomohiro Usui¹, Nathaniel T. Stein², Thomas Prettyman⁷, Andrea Raponi⁵, Mauro Ciarniello⁵ (1. Earth-Life Science Institute, Tokyo Institute of Technology, 2. California Institute of Technology, 3. Jet Propulsion Laboratory, 4. Agenzia Spaziale Italiana, 5. Institute for Space Astrophysics and Planetology, National Institute for Astrophysics, 6. Harvard University, 7. Planetary Science Institute)
Keywords: Ceres, Planetary exploration, Mineralogy

Quantitative determination of the mineral assemblages on the surface is important for the understanding of aqueous alteration on Ceres. Unveiling the aqueous paleoenvironment of Ceres would constrain the formation and evolution processes of Ceres, which is tied to the dynamical history of the early solar system. Infrared spectroscopic observations of Ceres from Dawn/VIR have shown that its dark surface contains Mg-phyllsilicates, ammonium-bearing phases, Mg/Ca carbonates, and dark materials, considered to be organic carbon and/or magnetite. The amount of a given material type present is currently constrained by “best-fits” of spectra with radiative transfer models using optical constants from terrestrial libraries [1].

In order to determine statistically rigorous error bars on quantification of material types present, we applied a Bayesian method to spectral unmixing of Ceres average surface. The Bayesian Markov Chain Monte Carlo method samples the posterior probability distribution function (PDF), i.e. the probability of a given set of mineral abundances and their grain sizes. This approach enables us to derive the mineral abundances on Ceres and the statistical error bars, taking observational errors in the reflectance spectra into account. As previous works [e.g., 1], we utilize the Hapke unmixing model for radiative transfer, which assumes the single-scattering albedo (SSA) spectrum of a surface is the sum of the SSA spectra of the material types comprising the surface, weighted by their cross sectional areas and proportional to abundance. Inputs are optical constants of the end member material types (n, k) as well as abundances range (0-100%) and grain sizes (10 μm -1 mm).

We obtained the best-fit composition of Ceres average surface which is consistent with the previous work utilised VIR data [1]. The Hapke model predicted the large amount of dark materials (proposed to be magnetite and/or amorphous carbon) up to ~80 wt.%, which is inconsistent with the elemental abundances estimated from the Gamma Ray and Neutron Detector (GRaND) [2]. We discuss the possible

agents of the darkening of Ceres surface.

[1] De Sanctis et al., 2015, Nature, 528, 241-244. [2] Prettyman T. H. et al., 2017, Science, 355, 6320, 55-59.