
 [JJ] Evening Poster | M (Multidisciplinary and Interdisciplinary) | M-TT Technology & Techniques

[M-TT37]Frontiers in Geochemistry

convener:Urumu Tsunogai(Graduate School of Environmental Studies, Nagoya University), Yoshio Takahashi(Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo), Tsuyoshi Iizuka(東京大学)

Sun. May 20, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Many new findings in earth and planetary sciences have been obtained by using state-of-the-art techniques supported by new technical development in analytical chemistry. This session aims at providing an opportunity for those developing new analytical methods to get together and have a strategic discussion on frontiers in geochemistry and cosmochemistry. We welcome a wide range of cutting-edge geochemical topics based on technical development, which have a potential for breakthrough of earth and planetary sciences. Besides, topics related to the direction of geochemistry and cosmochemistry in future are also welcome. Especially, we welcome topics which present how to install/maintain precious facilities in geochemical laboratories. We expect wide-ranged and future-oriented discussion to develop geochemistry and cosmochemistry.

[MTT37-P04]Organic complexation of zinc at the shallow hydrothermally active area of the Tachibana Bay, Nagasaki

Taejin Kim¹, *Hajime Obata¹, Asami S Mashio², Toshitaka Gamo¹, Shigenobu Takeda³ (1.Marine inorganic chemistry division, Atmosphere and Ocean Research Institute, The University of Tokyo, 2.College of Science and Engineering, School of Chemistry, Kanazawa University, 3.Graduate School of Fisheries Science and Environmental Science, Nagasaki University)

Keywords:Zinc, organic ligand, hydrothermal, sulfide

I. Introduction

Trace metals, like Zn, are used in numerous enzyme systems involved with a variety of metabolic processes. Low Zn concentrations in surface seawater could limit growth of some kinds of phytoplankton (Brand et al., 1983; Sunda and Huntsman, 2000). Therefore, it is an important subject to reveal the biogeochemical cycles of Zn in the ocean.

Recently, dissolved Zn enrichments near the hydrothermal vents have also been observed in the Mid-Atlantic Ridge and East Pacific Rise (Conway and John, 2014; Roshan et al., 2016), which suggested mantle-derived Zn from the hydrothermal vents could be a major source of dissolved Zn in ocean environment (Roshan et al., 2016). Sander and Koschinsky (2011) mentioned that organic ligands could potentially stabilize dissolved Zn in hydrothermal fluids and plumes. However, this process in the hydrothermal system has not been investigated enough yet.

In this study, we determined dissolved trace metal (Mn, Fe, Cu, and Zn) concentrations, Zn organic speciation, and dissolved sulfide in seawater in Tachibana Bay near the coast of Obama hot springs.

II. Sampling and Methods

Seawater samples were collected using acid-cleaned samplers with external springs mounted on CTD-CMS system during a research cruise KY-340 of TS *Kakuyo-maru*, May 2012. The samples were collected in low-density polyethylene bottles through a 0.2 μm -pore size filter. Samples for dissolved trace metal analysis were acidified to a pH of less than 1.8 using ultrapure HCl, and stored. Another set of samples, for Zn speciation analysis, was frozen at -18°C immediately after sampling. The samples were brought back to the laboratory and analyzed using CLE-CSV for Zn speciation (Kim et al., 2015), and using ICP-MS for trace metal concentrations (Kondo et al., 2016).

III. Results and Discussion

The ranges of dissolved Mn (13.3 – 39.3 nM), dissolved Fe (1.8 – 16.5 nM), and dissolved Cu (1.8 – 2.6 nM) in seawaters of Tachibana Bay were within the concentration levels in the East China Sea. The dissolved Zn concentrations ranged from 0.3 to 3.1 nM, which were also similar concentration level to those in the East China Sea.

Vertical distributions of dissolved trace metals (Mn, Fe, Cu, and Zn) at the coastal hydrothermal area showed different features compared to those at the center of Tachibana Bay. At the coastal area, dissolved Mn and Fe concentrations were generally high and increased toward the bottom, whereas dissolved Cu and Zn concentrations were decreased. On the other hand, vertical distributions of trace metals at the center of the bay were relatively constant. We discuss the role of organic complexing ligands and sulfide on Zn and Cu distributions in the coastal hydrothermal environment.

References

- Brand, L. E., W. G. Sunda, and R. R. Guillard (1983), Limitation of marine phytoplankton reproductive rates by zinc, manganese, and iron. *Limnology and Oceanography* 28, 1182–1198.
- Conway, T. M. and S. G. John (2014), The biogeochemical cycling of zinc and zinc isotopes in the North Atlantic Ocean. *Global Biogeochemical Cycles*, 28, 1111–1128.
- Kim, T., H. Obata, Y. Kondo, H. Ogawa, and T. Gamo (2015b), Distribution and speciation of dissolved zinc in the western North Pacific and its adjacent seas, *Marine Chemistry*, 173, 330–341.
- Kondo, Y., H. Obata, N. Hioki, A. Ooki, S. Nishino, T. Kikuchi, K. Kuma (2016), Transport of trace metals (Mn, Fe, Ni, Zn and Cd) in the western Arctic Ocean (Chukchi Sea and Canada Basin) in late summer 2012. *Deep-Sea Research-I*, 116, 236–252.
- Roshan, S., J. Wu, and W. J. Jenkins (2016), Long-range transport of hydrothermal dissolved Zn in the tropical South Pacific, *Marine Chemistry*, 183, 25–32.
- Sander, S. G. and A. Koschinsky (2011), Metal flux from hydrothermal vents increased by organic complexation. *Nature Geoscience*, 4, 145–150.

Sunda, W. G., and S. A. Huntsman (2000), Effect of Zn, Mn, and Fe on Cd accumulation in phytoplankton: Implications for oceanic Cd cycling. *Limnology and Oceanography*, 45, 1501–1516.