

---

[JJ] Evening Poster | M (Multidisciplinary and Interdisciplinary) | M-ZZ Others

## [M-ZZ41] Marine manganese deposits: from basic to applied sciences

convener: Akira Usui (Marine Core Research Center, Kochi University), Yoshio Takahashi (Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo), Katsuhiko Suzuki (国立研究開発法人海洋研究開発機構・海底資源研究開発センター, 共同), Takashi Ito (Faculty of Education, Ibaraki University)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Marine manganese deposits include nodules and crusts of massive iron-manganese oxide aggregates associated with useful metals. The deposits are known as potential resources of cobalt, copper, nickel, rare earth elements, platinum, and tellurium. However, the controlling parameters on the regional and temporal variations in chemical and mineralogical composition have not been clarified yet. In this session, various factors in the growth of manganese oxides, enrichment and circulation of metals, paleoenvironment, and formation age of manganese deposits will be discussed from viewpoints of geology, mineralogy, paleocean sciences, geochemistry, microbiology, and sea floor engineering.

---

## [MZZ41-P04] Vertical profile of neodymium isotopic composition in the oceanic region close to the Takuyo-Daigo Seamount

\*Hiroshi Amakawa<sup>1</sup>, Yusuke Fukami<sup>1</sup>, Junji Torimoto<sup>1</sup>, Tatsuo Nozaki<sup>2</sup>, Koichi Iijima<sup>2</sup>, Akira Usui<sup>3</sup>, Katsuhiko Suzuki<sup>2</sup> (1. Japan Agency for Marine-Earth Science Technology Research and Development (R&D) Center for Submarine Resources, 2. Research and Development Center for Submarine Resources, Japan Agency for Marine-Earth Science and Technology, 3. Marine Core Research Center, Kochi University)

Keywords: ferromanganese crust, Nd isotopic composition, seawater, Takuyo-Daigo Seamount, remotely operated vehicle

Ferromanganese crusts, which are chemical precipitates occurring on the ocean floor and seamount, are known to be rich in lanthanides including neodymium (Nd). The major source of lanthanides in the surface layer ferromanganese crust is ambient seawater. However, whether the lanthanides in ferromanganese crusts are supplied from ambient seawater or from the whole water column above ferromanganese crusts were not clearly presented. To identify the source of lanthanides (Nd) in the surface layer of ferromanganese crusts, the vertical profile of seawater Nd isotopic composition was determined for a location near the Takuyo-Daigo Seamount (21°59.2′N, 153°56.0′E) in the northwest Pacific Ocean. The data were compared with those of the surface layer ferromanganese crust obtained from the Takuyo-Daigo Seamount by remotely operated vehicles (ROVs), whose sampling depths ranged from 965 m to 5385 m. The seawater Nd isotopic composition profile was similar to the surface layer Nd isotopic composition of ferromanganese crusts within analytical errors. This confirms that Nd in the surface layer of ferromanganese crusts is supplied directly by ambient seawater.