

---

 [JJ] Evening Poster | P (Space and Planetary Sciences) | P-CG Complex & General

## [P-CG23] Planetary Magnetosphere, Ionosphere, and Atmosphere

convener: Kanako Seki (Graduate School of Science, University of Tokyo), Takeshi Imamura (Graduate School of Frontier Sciences, The University of Tokyo), Naoki Terada (東北大学大学院理学研究科, 共同), Hiroyuki Maezawa (Department of Physical Science Osaka Prefecture University)

Sun. May 20, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Exploration of Moon, Venus, Mars, Mercury, Jupiter, Saturn, and beyond together with rapid developments of numerical simulations provides us new view of planetary environment. This session collects general contributions of new findings about planetary magnetosphere, ionosphere, and atmosphere. New methodology and technology development studies for future explorations are also welcome. In order to put the common knowledge at different planets into perspective, this session aims to facilitate discussions on comparative planetary environments.

---

## [PCG23-P05] Full PIC simulation analysis on the plasma dynamics in a small-scale magnetosphere

\*Satoki Oki<sup>1</sup>, Hideyuki Usui<sup>1</sup>, Naoki Terada<sup>2</sup>, Kanako Seki<sup>3</sup>, Yohei Miyake<sup>1</sup>, Yuto Katoh<sup>2</sup>, Manabu Yagi<sup>4</sup>

(1. Graduate School of System Informatics, Kobe University, 2. Graduate School of Science, Tohoku University, 3. Graduate School of Science, University of Tokyo, 4. RIKEN Advanced Institute for Computational Science)

Keywords: Particle-in-cell simulation, Weakly magnetized airless body, small-scale magnetosphere, plasma dynamics

The objectives of this research is to study a small magnetosphere formed by the interaction between a weakly magnetized small body and the solar wind and to understand plasma dynamics at the orbit level of the particle by performing three-dimensional particle simulation. As examined in the previous works, Mercury has weak intrinsic magnetic field and a small-scale magnetosphere is created due to the interaction with the solar wind. The previous spacecraft observation revealed that the dipole moment of the Mercury's intrinsic magnetic field is around 1/2000 of that of the Earth and it causes a small (i.e. 1/20 of that of the Earth) magnetosphere. The proportion occupied by the Mercury body in the magnetosphere is large. The Mercury magnetosphere has been studied by MHD and hybrid particle simulation so far. However, plasma kinetics have not much considered in those previous simulations. In this study, we performed three-dimensional full particle simulations to understand the plasma dynamics including particle trajectories in the small magnetosphere. In the simulation model, the weakly magnetized small body is set in the center, and the solar wind having the southward IMF is injected from the boundary of the simulation domain. When we define  $D_p$  as the distance between the dipole center and a position where the solar wind dynamics pressure balances the magnetic pressure at the dayside, the ratio of  $D_p$  to ion inertia length  $\lambda_{Di}$  can be more than 10 in the Mercury model. In the present study, in order to emphasize the kinetic effect, we set  $\lambda_{Di}/D_p = 1$ . When we define  $R_b$  as the radius of sphere, we took a relatively large proportion of sphere in the magnetosphere as  $R_b/D_p = 0.6$ . We reported that dawn-dusk asymmetry characteristics of the plasma spatial distribution are found in the equatorial plane. In the present study, we examined the formation of the asymmetry distribution by considering the plasma dynamics including analysis about particle orbit. We particularly focused on the electron acceleration in the region where the southward IMF and the magnetospheric magnetic field cancel each other and the magnetic field intensity becomes zero. We also focused on the plasma dynamics in the cusp and the tail region of the magnetosphere and analyzed how the plasma particles

behave in the inner magnetosphere near small body. In addition, we investigated surface charging and electric field by considering the amount of plasma flux flowing into the surface of the body. The knowledge is important for understanding the distribution of ion ( $\text{Na}^+$ ) and its acceleration in the Mercury magnetosphere. In the future research, we will consider the contribution of heavy ions and photo-electrons with a certain scale height, and analyze their dynamics in the Mercury environment.