
[EE] Evening Poster | P (Space and Planetary Sciences) | P-EM Solar-Terrestrial Sciences, Space Electromagnetism & Space Environment

[P-EM10]Coupling Processes in the Atmosphere-Ionosphere System

convener:Huixin Liu(Earth and Planetary Science Division, Kyushu University SERC, Kyushu University), Loren Chang(Institute of Space Science, National Central University), Yuichi Otsuka(名古屋大学宇宙地球環境研究所)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Vertical coupling mechanisms throughout the whole atmosphere are critical to understanding the near Earth space environment, as well as its sensitivity to the solar, geomagnetic, and atmospheric drivers. This international session focuses on physical/chemical processes occurring in the mesosphere, thermosphere, and ionosphere (MTI) from both the poles to the equatorial region. Both quiet and disturbed states in response to lower atmospheric forcing or solar forcing are important for understanding the MTI system and its coupling to other regions. We invite presentations of observations and observational concepts with ground-based and/or space-borne instruments, theoretical studies, numerical simulations, and development of data analysis systems for various kinds of temporal and spatial variations in MTI system.

[PEM10-P04]Initial report on PMC observations by Himawari-8

*Takuo T. Tsuda¹, Yuta Hozumi¹, Kento Kawaura¹, Hidehiko Suzuki², Keisuke Hosokawa¹, Takuji Nakamura³ (1.University of Electro-Communications, 2.Meiji University, 3.National Institute of Polar Research)

Keywords:PMC, NLC, Himawari-8, Geostationary-Earth-Orbit satellite, Space-borne imaging

Polar mesospheric clouds (PMCs) or noctilucent clouds (NLCs) consist of water-ice particles, which can be produced at cold summer mesopause region, mainly at high latitudes. PMC observations have been widely performed by various methods from the ground as well as from the space. Optical observations by ground-based cameras, imagers or lidars are often limited by the weather, because the clear sky is required for such observations. Hence, satellite observations from the space are valuable for more continuous observations, which enable more systematic data coverage. Such systematic data coverage would be of benefit, for example, to monitor long-term PMC activities, which may be related with the global change because mesosphere cooling, which can enhance water-ice particle production, may be induced by CO₂ and CH₄ increases.

In this presentation, we will make an initial report on PMCs observed by Himawari-8, the Japanese Geostationary-Earth-Orbit (GEO) meteorological satellite. In the regular operation of Himawari-8, full-disk images of the Earth are obtained every 10 minutes with ~1 km spatial resolution. We can find PMC emissions in the Earth's limb region of the full-disk images. Thus, Himawari-8 provides PMC observations by continuous limb-viewing from its almost fixed locations relative to the Earth. We will introduce our initial results on Himawari-8 observed PMCs and discuss future PMC research utilizing the Himawari-8 observations.