

[EE] Evening Poster | P (Space and Planetary Sciences) | P-EM Solar-Terrestrial Sciences, Space Electromagnetism & Space Environment

[P-EM10]Coupling Processes in the Atmosphere-Ionosphere System

convener:Huixin Liu(Earth and Planetary Science Division, Kyushu University SERC, Kyushu University), Loren Chang(Institute of Space Science, National Central University), Yuichi Otsuka(名古屋大学宇宙地球環境研究所)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Vertical coupling mechanisms throughout the whole atmosphere are critical to understanding the near Earth space environment, as well as its sensitivity to the solar, geomagnetic, and atmospheric drivers. This international session focuses on physical/chemical processes occurring in the mesosphere, thermosphere, and ionosphere (MTI) from both the poles to the equatorial region. Both quiet and disturbed states in response to lower atmospheric forcing or solar forcing are important for understanding the MTI system and its coupling to other regions. We invite presentations of observations and observational concepts with ground-based and/or space-borne instruments, theoretical studies, numerical simulations, and development of data analysis systems for various kinds of temporal and spatial variations in MTI system.

[PEM10-P10]Effects of horizontal wind structure on the gravity wave activity in the upper stratosphere over Syowa

*Masaru Kogure^{1,2}, Takuji Nakamura^{2,1}, Mitsumu K. Ejiri^{2,1}, Takanori Nishiyama^{2,1}, Yoshihiro Tomikawa^{2,1}, Masaki Tsutsumi^{2,1} (1.The Graduate University for Advanced Studies, 2.National Institute of Polar Research)

Keywords:Gravity wave, Antarctic, Polar night jet, Lidar, Intermittency, middle atmosphere

Gravity waves (GWs) transport their momentum and energy from the lower atmosphere to the upper atmosphere and drive the general circulation, which significantly changes the temperature in the middle atmosphere [Fritts and Alexander, 2003]. To understand this role quantitatively will improve the modern general circulation models. Typical GW activities and its seasonal variations have been studied at various places around the world. Some studies have shown global maps of the typical activities from satellite observations lately. However, what causes a shorter temporal and local variations of the activities are poorly understood, especially above the upper stratosphere.

To understand this cause, we estimated the potential energy of GW (E_p) over Syowa Station (69°S, 40°E) from a Rayleigh/Raman (RR) lidar observation between 2011 and 2015. We found a clear enhancement of the E_p during 8th-21st August 2014. The E_p in this period was about two and five times as large as the winter mean in the other years at 50 and 60 km, while the one between 20 and 40 km was as small as the winter mean. One of possibility is that some source would exist around 45 km, but there seemed no specific source within the Modern-Era Retrospective analysis for Research and Applications (MERRA) [Rienecker et al., 2011]. The second one is that the GWs would converge from the lower atmosphere at various latitudes due to the meridional gradient of the westerly wind [Dunkerton, 1984]. The polar night jet around 40°E during the enhancement period slanted to ~70°S from ~50°S and the gradient was larger. In such condition, the GWs with west-ward wavenumber can converge to Syowa. We estimated paths of GWs in order to examine whether the GWs could converge to Syowa. We found that the large-scale GWs could converge to 50-55 km altitudes over Syowa in that

period. This result suggests that the enhancement could be caused by the convergence of the GWs.

In this presentation, we will show the potential energy, the wind field and estimated paths of the GWs during the enhancement period, and discuss the cause of the enhancement.