

[EE] Evening Poster | P (Space and Planetary Sciences) | P-EM Solar-Terrestrial Sciences, Space Electromagnetism & Space Environment

[P-EM16] Dynamics of Earth's Inner Magnetosphere and Initial Results from Arase

convener: Danny Summers (Memorial University of Newfoundland), Yoshizumi Miyoshi (Institute for Space-Earth Environmental Research, Nagoya University), Keisuke Hosokawa (電気通信大学大学院情報理工学研究科, 共同), Yusuke Ebihara (Research Institute for Sustainable Humanosphere, Kyoto University)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Earth's inner magnetosphere is a fascinating source of space research problems. There remain many fundamental questions concerning the physics of the radiation belts, the ring current, the plasmasphere and the ionosphere. The JAXA spacecraft Arase (ERG) was successfully launched in December 2016, and has since been providing excellent data on waves, particles and fields over a range of L-shells in the inner magnetosphere. This session particularly welcome submissions related to the Arase mission. As well, data from other recent missions to the magnetosphere are also welcome, including the Van Allen Probes, MMS, and THEMIS. Topics of interest include charged particle interactions with the predominant electromagnetic wave modes such as whistler-mode chorus and hiss, ion cyclotron waves, magnetosonic waves, and ULF waves. Projects involving the prevailing issues of particle acceleration and loss, and particle transport are also of interest. In addition, projects involving the coupling of plasma populations in the inner magnetosphere are also timely. Studies involving observations, simulations, theory and modeling are all invited.

[PEM16-P10] Model Coupling between BATS-R-US - CIMI and GEMSIS-RC for study of ULF waves in the inner magnetosphere

*Kei Kamiya¹, Kanako Seki², Shinji Saito¹, Takanobu Amano², Yoshizumi Miyoshi³, Mei-Ching Fok⁴, Colin Komar⁴ (1. Graduate School of Science, Nagoya University, 2. Graduate School of Science, University of Tokyo, 3. Institute for Space-Earth Environmental Research, Nagoya University, 4. Goddard Space Flight Center, NASA)

Keywords: ULF waves in the inner magnetosphere, Model Coupling between MHD and Ring Current

Radiation belts are regions of enhanced population of relativistic electrons with energies over MeV in the Earth's inner magnetosphere. The outer radiation belt electrons exhibit significant variations during geomagnetic storms. There are two major candidates of the acceleration mechanism: the radial transport from external source and local acceleration within the inner magnetosphere. Previous observations have used radial profiles of the phase space density (PSD) to distinguish two mechanisms [e.g. Reeves et al., Science, 2013]. While the local acceleration creates a localized peak in the PSD radial profile, the radial transport has been considered as a diffusive process that reduces gradient in the PSD profile. However, under certain conditions, the radial transport process can also produce localized peaks in the PSD profile [e.g. Degeling et al., JGR, 2008]. Therefore, it is important to have another observable measure to distinguish the two acceleration mechanisms to understand the relativistic electron variations in the outer radiation belt.

Pc5 Ultra-Low Frequency (ULF) waves in the inner magnetosphere are observed as electromagnetic fluctuations with frequencies of 1.67-6.67 mHz and considered as a driver of the radial transport of relativistic electrons in the outer radiation belt. In our previous study, we showed that interaction with continuous monochromatic Pc5 waves can form characteristic butterfly pitch angle distributions of the

relativistic electrons in specific regions in the outer radiation belt determined by the drift-resonance condition [Kamiya et al., submitted to JGR, 2018]. This result suggests a possibility that the characteristic pitch angle distributions can be a measure to distinguish the radial transport driven by the Pc5 waves. On the other hand, the formation of the butterfly pitch angle distributions depends on the latitudinal distribution of the wave power and spectrum of the Pc5 waves. In order to model more realistic spatial distributions of the ULF waves for investigation of its interaction with relativistic electrons, we need to improve the outer boundary conditions in our global simulation of the inner magnetosphere to more realistic ones.

For this purpose, we have developed a model coupling method between two models, i.e., Global MHD-based simulation of the magnetosphere, Brock-Adaptive-Tree Solar-Wind Roe-Type Upwind Scheme & Comprehensive Inner Magnetosphere and Ionosphere (BATS-R-US - CIMI) and a 5-D drift-kinetic global inner magnetospheric simulation, GEMSIS-Ring Current (RC). The BATS-R-US - CIMI model solves the ideal MHD equations for the regions from the sunward boundary to the inner boundary at $L=2.5$ including the ring current pressure gradient [Glocer et al., JGR, 2013; Fok et al., JGR, 2014]. The GEMSIS-RC model solves the ring current ion PSD in the 5-D phase space together with time evolution of the electromagnetic fields self-consistently, which include the ULF waves in the inner magnetosphere ($3.6 < L < 6.6$) by calculating the drift-kinetic equations and Maxwell equations simultaneously [Amano et al., JGR, 2011]. We adopted time-dependent boundary between two models, which is located on the last closed field line in the GEMSIS-RC simulation domain. In the presentation, we will report on the validity assessment of the model coupling method and the initial result of the ULF wave simulation in the inner magnetosphere with realistic boundary conditions.

References:

Reeves et al. (2013), "Electron Acceleration in the Heart of the Van Allen Radiation Belts", Science, 341, 30 AUGUST.

Degeling et al. (2008), "Drift resonant generation of peaked relativistic electron distributions by Pc 5 ULF waves", JGR, 113, A02208, doi:10.1029/2007JA012411.

Kamiya et al. (Submitted in 2018), "Formation of butterfly pitch angle distributions of relativistic electrons in the outer radiation belt with a monochromatic Pc5 wave", JGR

Glocer et al. (2013), "CRCM + BATS-R-US two-way coupling", JGR, 118, 1635–1650, doi:10.1002/jgra.50221

Fok et al. (2014), "The Comprehensive Inner Magnetosphere-Ionosphere Model", JGR, 119, 7522–7540, doi:10.1002/2014JA020239.

Amano et al. (2011), "Self-consistent kinetic numerical simulation model for ring current particles in the Earth's inner magnetosphere", JGR, 116, A02216, doi:10.1029/2010JA015682.