

[EE] Evening Poster | P (Space and Planetary Sciences) | P-EM Solar-Terrestrial Sciences, Space Electromagnetism & Space Environment

[P-EM16] Dynamics of Earth's Inner Magnetosphere and Initial Results from Arase

convener: Danny Summers (Memorial University of Newfoundland), Yoshizumi Miyoshi (Institute for Space-Earth Environmental Research, Nagoya University), Keisuke Hosokawa (電気通信大学大学院情報理工学研究科, 共同), Yusuke Ebihara (Research Institute for Sustainable Humanosphere, Kyoto University)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Earth's inner magnetosphere is a fascinating source of space research problems. There remain many fundamental questions concerning the physics of the radiation belts, the ring current, the plasmasphere and the ionosphere. The JAXA spacecraft Arase (ERG) was successfully launched in December 2016, and has since been providing excellent data on waves, particles and fields over a range of L-shells in the inner magnetosphere. This session particularly welcome submissions related to the Arase mission. As well, data from other recent missions to the magnetosphere are also welcome, including the Van Allen Probes, MMS, and THEMIS. Topics of interest include charged particle interactions with the predominant electromagnetic wave modes such as whistler-mode chorus and hiss, ion cyclotron waves, magnetosonic waves, and ULF waves. Projects involving the prevailing issues of particle acceleration and loss, and particle transport are also of interest. In addition, projects involving the coupling of plasma populations in the inner magnetosphere are also timely. Studies involving observations, simulations, theory and modeling are all invited.

[PEM16-P12] Temporal and spatial correspondence of EMIC waves and energetic electron precipitation observed by ground-based stations on 27 March 2017

*Hirai Asuka¹, Fuminori Tsuchiya¹, Takahiro Obara¹, Hiroaki Misawa¹, Kazuo Shiokawa², Yoshizumi Miyoshi², Martin G Connors³, Satoshi Kurita² (1. Planetary Plasma and Atmospheric Research Center, Graduate School of Science, Tohoku Univ., 2. Institute for Space-Earth Environmental Research, Nagoya University, 3. Athabasca University)

Keywords: EMIC waves, energetic electron precipitation

Energetic electron losses from the outer radiation belt occur during magnetic storms and substorms. One of the mechanisms is precipitation of the electrons into the atmosphere. The electromagnetic ion cyclotron (EMIC) wave is one of the candidates to cause pitch angle scattering of the energetic electrons. EMIC waves, which are observed in the Pc1–Pc2 frequency range (0.1–5 Hz), are excited by the ion cyclotron instability in the equatorial region of the magnetosphere and propagate along magnetic field lines to the ionosphere. While it has been theoretically studied that EMIC waves play an important role in the energetic electron precipitation, it is not easy to observe temporal and spatial proximities of these phenomena.

Here, we investigate relationship between occurrence of EMIC waves and energetic electron precipitation using multiple ground-based observations.

We used induction magnetometer data in North America (PWING and CARISMA stations) to find occurrence of EMIC waves. We also use artificial VLF/LF radio waves that propagate from transmitters to a receiver through reflection, between earth's surface and the lower ionospheric boundary at altitudes of 70–90 km. Ionization caused by precipitating electrons on the radio propagation path results in changes in the propagation path and attenuation of the signal amplitude. Thus, precipitation of energetic electrons at

energies higher than $\sim 100\text{keV}$ causes deviation of the VLF/LF wave amplitude from that in undisturbed conditions. In this study, we analyzed VLF/LF wave signals received at Athabasca, Canada (latitude=54.7, longitude=246.7, L=4.45). An all sky imager at Athabasca (one of the PWING stations) was used to observe spatial and temporal characteristics of proton aurora which was useful to identify ionospheric sources of the EMIC waves.

During the main phase of a geomagnetic storm on 27 March 2017, Pc1 waves were observed at several stations: Kapuskasing (latitude=49.4, longitude=277.81, L=4.72), Pinawa (latitude=50.2, longitude=263.7, L=4.06), Thief River Falls (latitude=48.0, longitude=263.6, L=3.58), Ministik Lake (latitude=53.4 longitude=247.0, L=4.22) and Athabasca in 03:10-06:30 UT. Pc1 waves were initially seen at Kapuskasing and the observed locations moved westward. Simultaneously, energetic electron precipitations were detected by VLF/LF radio waves transmitted from several stations in the United States: WWVB (latitude=40.7, longitude=255.0, L=2.28, 60kHz), NDK (latitude=46.4, longitude=261.5, L=2.98, 25.2kHz) and NLK (latitude=48.2, Longitude=238.1, L=2.85, 24.79kHz).

Isolated proton auroras appeared on the VLF/LF radio propagation paths when energetic electron precipitation was detected. When the proton aurora propagated westward and departed from the radio propagation paths, the energetic electron precipitation ceased.

The data sets presented here indicate close temporal and spatial relationship between EMIC waves and energetic electron precipitation and support that the EMIC waves drive precipitation of energetic electrons and protons into the atmosphere.

Acknowledgement

The authors thank I.R. Mann, D.K. Milling and the rest of the CARISMA team for data. CARISMA is operated by the University of Alberta, funded by the Canadian Space Agency.