

[EE] Evening Poster | P (Space and Planetary Sciences) | P-EM Solar-Terrestrial Sciences, Space Electromagnetism & Space Environment

[P-EM16] Dynamics of Earth's Inner Magnetosphere and Initial Results from Arase

convener: Danny Summers (Memorial University of Newfoundland), Yoshizumi Miyoshi (Institute for Space-Earth Environmental Research, Nagoya University), Keisuke Hosokawa (電気通信大学大学院情報理工学研究科, 共同), Yusuke Ebihara (Research Institute for Sustainable Humanosphere, Kyoto University)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Earth's inner magnetosphere is a fascinating source of space research problems. There remain many fundamental questions concerning the physics of the radiation belts, the ring current, the plasmasphere and the ionosphere. The JAXA spacecraft Arase (ERG) was successfully launched in December 2016, and has since been providing excellent data on waves, particles and fields over a range of L-shells in the inner magnetosphere. This session particularly welcome submissions related to the Arase mission. As well, data from other recent missions to the magnetosphere are also welcome, including the Van Allen Probes, MMS, and THEMIS. Topics of interest include charged particle interactions with the predominant electromagnetic wave modes such as whistler-mode chorus and hiss, ion cyclotron waves, magnetosonic waves, and ULF waves. Projects involving the prevailing issues of particle acceleration and loss, and particle transport are also of interest. In addition, projects involving the coupling of plasma populations in the inner magnetosphere are also timely. Studies involving observations, simulations, theory and modeling are all invited.

[PEM16-P14] Excitation mechanism of storm-time Pc5 ULF waves by ring current ions

*Tomotsugu Yamakawa¹, Kanako Seki¹, Takanobu Amano¹, Kunihiro Keika¹, Yoshizumi Miyoshi², Naoko Takahashi¹ (1. School of Science, The University of Tokyo, 2. ISEE, Nagoya University)

Keywords: ring current, magnetic storm, Pc5 ULF, drift-bounce resonance, drift-kinetic simulation

Storm-time Pc5 ULF waves have been considered to be generated by ring current particles associated with plasma injection into the inner magnetosphere during magnetic storms. It is generally believed that ULF waves drive radial diffusion of radiation belt relativistic electrons (e.g., Shprits et al., 2008). Therefore, it is important to know how Pc5 ULF waves are generated by ring current ions to understand the acceleration mechanism of radiation belt electrons. One primary excitation mechanism of storm-time Pc5 waves is the resonance occurring between the drifting and bouncing motions of ring current ions (Southwood, 1976). Although the drift-bounce resonance excitation of these waves is shown by many observations (e.g., Ozeke and Mann, 2008), fundamental features of the excitation mechanism such as the growth rate and necessary phase space density (PSD) distributions is far from understood. One of the reason is due to difficulty of non-linear global simulations of the excitation mechanisms.

In order to simulate the excitation mechanism of the storm-time Pc5 waves, we perform a kinetic simulation for ring current particles using GEMSIS-RC model (Amano et al., 2011), in which five-dimensional drift-kinetic equation for PSD of ring current ions and Maxwell equation are solved self-consistently, and the first adiabatic invariant is assumed to be conserved. In order to simplify injection model, we set a localized high pressure region around midnight consisting of O⁺ ions which have a Maxwellian velocity distribution of the temperature 16 keV (O⁺ ion) with a loss cone. In addition, the presence of a background cold dense plasma, which is fixed to the initial condition is assumed. The fixed boundary condition for the field quantities is used, and the mirror reflection at the ionosphere is assumed at the inner boundary.

We investigate the power spectrum of magnetic field fluctuations, and it is shown that primarily poloidal mode (Br) is excited from duskside to afternoon magnetic local times. The frequency of the excited waves matches the expectation from the drift-bounce resonance conditions for the locally dominant O⁺ ions estimated from the simulated PSD. It is also found that poloidal mode wave propagates westward from 1800 to 1200 MLT associated with the westward transport of O⁺ ions. We also investigate the local PSD distribution, and the systematic modulation of the ion pitch angle distribution associated with the phase of excited waves are confirmed. Detailed correspondence of the PSD distributions and the wave properties will be also performed.

References

- Amano, T., K. Seki, Y. Miyoshi, T. Umeda, Y. Matsumoto, Y. Ebihara, and S. Saito (2011), Self-consistent kinetic numerical simulation model for ring current particles in the Earth's inner magnetosphere, *J. Geophys. Res.*, 116, A02216, doi:10.1029/2010JA015682.
- Ozeke, L. G., and I. R. Mann (2008), Energization of radiation belt electrons by ring current ion driven ULF waves, *Journal of Geophysical Research: Space Physics*, 113(A2), A02,201, doi:10.1029/2007JA012468.
- Shprits, Y. Y., S. R. Elkington, N. P. Meredith, and D. A. Subbotin (2008), Review of modeling of losses and sources of relativistic electrons in the outer radiation belt I: Radial transport, *J. Atmos. Sol. Terr. Phys.*, 70(14), 1679 – 1693, doi: 10.1016/j.jastp.2008.06.008.
- Southwood, D. J. (1976), A general approach to low-frequency instability in the ring current plasma, *Journal of Geophysical Research*, 81(19), 3340-3348, doi: 10.1029/JA081i019p03340.