[EE] Evening Poster | P (Space and Planetary Sciences) | P-PS Planetary Sciences

[P-PS01]Outer Solar System Exploration Today, and Tomorrow

convener:Jun Kimura(Osaka University), Yasumasa Kasaba(Dep. Geophysics Graduate School of Science Tohoku University), Steven Vance(Jet Propulsion Laboratory, Caltech, 共同), Kunio M. Sayanagi (Hampton University)

Mon. May 21, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe) The giant planets provide many keys to understanding planetary processes. They play an important role in shaping our solar system, and the physical and chemical processes they harbor also provide a unique opportunity to study the phenomena relevant for studying

Earth and other planets, including exoplanetary systems. In this session, we discuss a wide range of topics encompassing the giant planets and their moons, including their origins, interiors, atmospheres, compositions, surface features, and electromagnetic fields. To advocate for current and future outer planets exploration (Cassini, Juno, New Horizons, JUICE, and beyond), we also call for discussions on future missions to explore giant planet systems, including how to develop better international cooperation. Discussion in this latter category will include progress in developing a solar sail mission concept for observing the Jupiter system and its trojan asteroids.

[PPS01-P09]A simulation study on solar wind influence on long-term variation of Jovian Synchrotron Radiation

*Sooman Han¹, Go Murakami², Hajime Kita³, Fuminori Tsuchiya³, Chihiro Tao⁴, Daniel Santos-Costa⁵, Atsushi Yamazaki², Masato Nakamura² (1.Department of Earh and Planetary Science, Graduate School of Science, The University of Tokyo, 2.Institue of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3.Planetary Plasma and Atmospheric Research Center, Tohoku University, 4.National Institue of Information and Communications Technology, 5.Southwest Research Institute) Keywords:Hisaki, Jupiter, Synchrotron, Radiation, Solar

Jovian Synchrotron Radiation (JSR) is a non-thermal radiation emitted by relativistic electrons trapped by Jovian magnetic field, whose frequency range is usually between a few hundred MHz and a few GHz. Based on the assumption that the source of electron is at some outer boundary (e.g. 6 Jovian Radii) and radial diffusion of electrons occur due to the violation of third adiabatic invariant, a number of diffusion models in the past could explain the steady profile of electron population and the resulted total flux density of JSR. Yet, the reason behind its time variation remains unclear, especially, no model could reproduce the long-term variation whose JSR amplitude is 20 ~ 30%, which has a strong correlation with solar wind dynamic pressure shifted 2 ~ 3 years forward in time.

In our study, we have made a radial diffusion model taking Hisaki's observation result into account that dawn-to-dusk electric field is present and is modulated with solar wind dynamic pressure. New diffusion coefficient is applied here in terms of solar wind dynamic pressure, and relative JSR amplitude is calculated according to the pressure variation between 1971 and 2005. By comparing with an observation at 2.3 GHz, we present our result that JSR amplitude of 20 ~ 30 % and a strong correlation between JSR and time-shifted solar wind dynamic pressure can be successfully recovered between 1971 and 1989, and discuss why and how long-term variation happens. For the period later than 1989 where the correlation is mostly lost, one might have to look for another possibility (e.g. active volcanic lo's activity).