[EE] Evening Poster | P (Space and Planetary Sciences) | P-PS Planetary Sciences

[P-PS01]Outer Solar System Exploration Today, and Tomorrow

convener:Jun Kimura(Osaka University), Yasumasa Kasaba(Dep. Geophysics Graduate School of Science Tohoku University), Steven Vance(Jet Propulsion Laboratory, Caltech, 共同), Kunio M. Sayanagi (Hampton University)

Mon. May 21, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe) The giant planets provide many keys to understanding planetary processes. They play an important role in shaping our solar system, and the physical and chemical processes they harbor also provide a unique opportunity to study the phenomena relevant for studying

Earth and other planets, including exoplanetary systems. In this session, we discuss a wide range of topics encompassing the giant planets and their moons, including their origins, interiors, atmospheres, compositions, surface features, and electromagnetic fields. To advocate for current and future outer planets exploration (Cassini, Juno, New Horizons, JUICE, and beyond), we also call for discussions on future missions to explore giant planet systems, including how to develop better international cooperation. Discussion in this latter category will include progress in developing a solar sail mission concept for observing the Jupiter system and its trojan asteroids.

[PPS01-P15]Longitudinal variations of the sulfur ions in the lo plasma torus observed by the HISAKI/EXCEED

Arakawa Ryo¹, *Hiroaki Misawa¹, Fuminori Tsuchiya¹, Masato Kagitani¹, Ichiro Yoshikawa², Kazuo Yoshioka², Hikida Reina³, Fumiharu Suzuki², Tomoki Kimura⁴, Go Murakami⁵, Atsushi Yamazaki⁵ (1.Planetary Plasma and Atmospheric Research Center, Graduate School of Science, Tohoku University, 2.Graduate School of frontier Science, The University of Tokyo, 3.Graduate School of Science, The University of Tokyo, 4.Nishina-Center for Accelerator Based Science, RIKEN, 5.Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency) Keywords:Jupiter, Io, volcanic event, periodicity, HISAKI

Previous ground-based and probe observations of Io plasma torus (IPT) in various wavelengths have detected a periodic variation which is longer than the System III period (9.925 h). It has been called the System IV periodicity.

In previous studies, the time variations in the System III longitude at the peak of the longitudinal modulation of several ions compositions and also those in the amplitude of them were investigated with the data obtained by the Cassini spacecraft [Steffl et al., 2006]. They found the following three features of the longitudinal modulation of the IPT. (i) The peak longitude shifted with respect to the System III longitude at the rate of 12.5deg/day, which shows the appearance of the System IV periodicity. (ii) The amplitude varied with the beat between the System III and IV periods, which shows that the IPT has the two periodicities (System III and IV). (iii) The phase relation between the densities of S⁺ and S³⁺ was 180degout of phase. They proposed the longitudinal 1-D physical chemistry model named as “Dual hot electron model (DHE model)” to reproduce the results obtained by the Cassini spacecraft. This model includes two longitudinal modulations of hot electron. While one rotates with the System III period, another one sub-corotates with the System IV period. The System III periodicity could be produced by the longitudinal inhomogeneity of the Jovian magnetic field [Hess et al., 2011]. However, the origin of the System IV periodicity has been unknown.

In this study, we analyzed time variations in intensities of EUV emissions from the IPT obtained by the HISAKI satellite to understand the mechanism responsible for the System IV periodicity. The data used in

this study was obtained during "1st season” (19 Dec. 2013 – 24 Apr. 2015), “2nd season” (27 Nov. 2014 – 14 May 2015) and “3rd season” (21 Jan. – 30 Aug. 2016). During the 2nd season, the lo’s volcanic event from the middle of Jan. to Mar. 2015 was reported from the ground based observation of logenic sodium emission [Yoneda et al., 2015]. To find temporal variations in the System IV period, temporal variations in the System III longitude at the peak of EUV intensity were derived by the function fitting to the light curves of three ion species (S⁺ at 76.5 nm, S²⁺ at 68.0 nm, and S³⁺ at 65.7 nm+140.5 nm). The System IV periods of S⁺ before and after the lo’s volcanic event were 10.0461±0.0097 h and 9.9891±0.0084 h, respectively. On the other hands, the System IV period of S⁺ was 9.9370±0.014 h during the lo’s volcanic event. This result is the first observation evidence which shows that the System IV period has shortened after the lo volcanic event with the data for the period including the whole series of the lo volcanic event.

The timing when the System IV period became short roughly corresponded to the timing when transient aurora activities enhanced. The phase of the longitudinal modulation of S³⁺ also abruptly changed in the order from the outer to the inner at that timing. These facts show the existence of the injection. We examined the effect of the gradient B drift of the injected hot electron and the collision relaxation time scale of the injected hot electron with the thermal electron. As the result, we suggest that the short System IV period just after the injection was produced by the gradient B drift of the injected hot electron whose energy was a few keV & Andash; a few ten keV. However, the origin of the & &Idquo;normal” System IV period, that is the period before the lo volcanic event, is not still understood.

Moreover, we investigated the phase relation between the densities of S^+ and S^{3+} to examine the DHE model with the spectral diagnosis for the data obtained by the HISAKI satellite. As the result, we found that the phase relation between the densities of S^+ and S^{3+} was not always 180deg out of phase, which was not consistent with the DHE model. Thus, additional or new ideas would be needed to explain these complex longitudinal modulations.