
[EJ] Evening Poster | P (Space and Planetary Sciences) | P-PS Planetary Sciences

[P-PS05]Lunar science and exploration

convener:Hiroshi Nagaoka(Waseda Univ.), Tomokatsu Morota(Graduate School of Environmental Studies, Nagoya University), Masaki N (名古屋大学宇宙地球環境研究所, 共同), Masahiro KAYAMA(Department of Earth and Planetary Material Sciences, Faculty of Science, Tohoku University)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Scientific data sets acquired by not only Japanese lunar mission SELENE (Kaguya), but also other countries' missions, have become new standard for lunar science. Analyses of these data have been providing several new knowledge and changing some hypotheses into the truth of the Moon. In concurrence with these studies, some countries including Japan are planning future lunar missions. In this session, we will discuss scientific results based on newly acquired lunar data, strategy for future missions including SLIM, and theoretical and experimental studies for lunar science.

[PPS05-P09]A new model of the boundary layer of the lunar wake

*Masaki N Nishino¹, Yoshifumi Saito², Hideo Tsunakawa³, Yuki Harada⁴, Shoichiro Yokota⁵, Futoshi Takahashi⁶, Masaki Matsushima³, Hidetoshi Shibuya⁷, Hisayoshi Shimizu⁸ (1.ISEE, Nagoya University, 2.ISAS/JAXA, 3.Tokyo Institute of Technology, 4.University of Iowa, 5.Osaka University, 6.Kyushu University, 7.Kumamoto University, 8.ERI, The University of Tokyo)

Keywords:Lunar wake, Solar wind, Electric field, Kaguya (SELENE)

Refilling of the tenuous lunar wake by solar wind plasma has been one of the fundamental phenomena of planetary plasma sciences. Because a portion of the solar wind electrons has much higher speed than protons, it has been widely accepted that suprathermal electrons precede protons to come into the wake along the interplanetary magnetic field. In this model, ambipolar (inward) electric fields around the wake boundary generated by the charge separation attract the surrounding solar wind protons into the central lunar wake. However, such treatment has implicitly assumed one-dimensional motion of the solar wind plasma along the magnetic field. Here we propose a new model of the wake boundary close to the Moon, taking into account the three-dimensionality of solar wind proton motions; Solar wind protons come into the lunar wake owing to their gyro motion and large inertia without help of suprathermal electrons, and those protons form an ion boundary layer between the surrounding solar wind and the tenuous region deeper inside the wake. We call this layer the wake ion boundary layer (WIBL). In this model, ambipolar electric fields would exist at the inner edge of the WIBL. This new model well explains electron signatures around the wake boundary detected by Kaguya (SELENE) at ~100 km altitude from the lunar surface. In addition, we suggest that the potential drop in the WIBL along the Kaguya orbit is several tens V, which is an order-of-magnitude smaller than the values previously reported as the wake boundary potential (400 V by Wind observation and 480 V by NOZOMI). We will discuss the reason of this discrepancy to obtain a comprehensive view of the lunar wake boundary.