

[EJ] Evening Poster | P (Space and Planetary Sciences) | P-PS Planetary Sciences

[P-PS05]Lunar science and exploration

convener:Hiroshi Nagaoka(Waseda Univ.), Tomokatsu Morota(Graduate School of Environmental Studies, Nagoya University), Masaki N (名古屋大学宇宙地球環境研究所, 共同), Masahiro KAYAMA(Department of Earth and Planetary Material Sciences, Faculty of Science, Tohoku University)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Scientific data sets acquired by not only Japanese lunar mission SELENE (Kaguya), but also other countries' missions, have become new standard for lunar science. Analyses of these data have been providing several new knowledge and changing some hypotheses into the truth of the Moon. In concurrence with these studies, some countries including Japan are planning future lunar missions. In this session, we will discuss scientific results based on newly acquired lunar data, strategy for future missions including SLIM, and theoretical and experimental studies for lunar science.

[PPS05-P13]Yield strength and lava tube cave height estimated from pits and lava flows of the Moon and Mars

*Tsutomu Honda¹ (1.Vulcano-speleological Society)

Keywords:Moon, Mars, vertical pit, lava flow, yield strength, lava tube

[Introduction] The vertical pit, Marius Hills Hole (MHH) of the Moon, found by Haruyama et al^(1~3) has several lava layers in it's cross-section as reported by Robinson et al⁽⁴⁾ through LRO observation. The cross sectional thickness is also reported from Cushing⁽⁵⁾ on the vertical pit in Martian Arsia Mons. On the other hand, the lava flow thickness of the Moon and Mars have been observed from the surface appearance of the lava flow from which lava yield strength were estimated, though these yield strengths had widely spread values. Here, the yields strengths are estimated from the thickness of the lava layers in a vertical pit and compared with those estimated from the lava flow surface appearance thickness. The height of lava cave tube possibly located under the vertical pit of the Moon and the Mars will be estimated by using the proper value from the comparison.

[The yield strength estimated from lava layer in the cross section of the vertical pit] Marius Hills Hole (MHH) consists of 4m-12m thickness of stratified lava layer in a vertical pit section(an average of 6m thickness)(Robinson⁽⁴⁾). When average thickness of $H=6\text{m}$ and slope angle $\alpha=0.31\text{ deg}$ in Rille-A area(Greeley⁽⁶⁾) are used for the lava flow stop condition of Simple flow, the yield strength is given as $f_b = \rho g \sin \alpha$; $H=131\text{ Pa}$ (Honda⁽⁷⁾) where the lava density is $\rho=2.5\text{g/cm}^3$ and surface gravity is $g=162\text{ cm/s}^2$. On the other hand, the thickness of the stratified lava layer of the ceiling section of vertical pit-I at the foot of north area in Arsia Mons is found to be $H=3\text{m}$ (Cushing⁽⁵⁾). The stop condition of Simple flow of lava where slope angle of pit-I area is 0.28 deg gives the yield strength $f_b = 136\text{ Pa}$. For this estimation, the lava density $\rho=2.5\text{g/cm}^3$ and surface gravity $g=371\text{ cm/s}^2$ are used.

[The yield strength obtained from the surface appearance thickness of the lava flow and comparison] Lots of yield strength of lava were obtained for lava flows of the Moon and Mars by using Simple flow stop condition (Hulme⁽⁸⁾), though these values are widely scattered. The Table 1 and Table 2 show the minimum and maximum value for the Moon and the Mars ever obtained. The minimum value 100 Pa in the Moon is near the yield strength 131 Pa obtained from cross sectional layer thickness of the pit. When the yield strength shows a bigger values, it seems that a lava flow manifests Multiple flow or Inflation of lava instead of Simple flow, then, the yield strength is considered as an apparent yield strength. An

example of lava thickness and yield strength as a function of slope angle of Arsia Mons of the Mars obtained by Moore et al⁽⁹⁾ is shown in Fig1 and Fig.2. The yield strength indicates the smaller value for the lower slope angle. The minimum value 120 Pa is near 136 Pa which is obtained from cross sectional layer thickness of the pit. For the lower slope angle, it seems probably to converge into the true yield strength.

[Estimation of the lava tube cave height under the pit] The lava tube cave height H_c under the MMH and Pit-I of Martian Arsia Mons will be estimated by the lava flow model on the inclined surface with slope angle α . The flow critical condition of the lava is expressed as $H_c = n f_g / (\rho g \sin \alpha)$, where ρ is density, g is gravity, For the case of $n=2$, cave height between infinite width parallel plates and for the case of $n=4$, cave height in the circular tube (Hulme⁽⁸⁾). The used yield strengths as proper value would be those obtained from the lava layer in the pit hole. For the MHH, for $n=2$ and $n=4$, $H_c=12\text{m}$ and 24m respectively. As the observed H_c is 17m , n will be about 3. This possibly shows the cross section of the lava tube cave under MHH is rectangular (Honda⁽⁷⁾). For the pit-I of Martian Arsia Mons, for $n=2$ and $n=4$, $H_c=6\text{m}$ and 12m respectively.

[Summary] The cross sectional observation of the lava flow layer is very important (Hasenaka et al⁽¹⁰⁾) to obtain the true yield strength. Because it seems difficult to judge whether the lava flow is Simple flow, or Multiple flow or Inflated flow from the remote surface appearance of the lava flow for the Moon and Mars. If the yield strength obtained from cross sectional observation contains still an influence of lava inflation, on site sampling and examination is necessary to obtain the true yield strength. With true yield strength, the lava tube cave height can be estimated.

References: (1)~(10), see Japanese version