[EE] Evening Poster | S (Solid Earth Sciences) | S-CG Complex & General

[S-CG53]Science of slow earthquakes: Toward unified understandings of whole earthquake process

convener:Satoshi Ide(Department of Earth an Planetary Science, University of Tokyo), Hitoshi Hirose(Research Center for Urban Safety and Security, Kobe University), Kohtaro Ujiie(筑波大学生命環境 系, 共同), Takahiro Hatano(Earthquake Research Institute, University of Tokyo)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe) Accumulating observational studies on various types of slow deformation events, such as tectonic tremors, very low frequency events, and slow slip events, portrays some universal characteristics in generally complex behavior, including interaction among events and influence by various outer loadings. Some of these phenomena seem to have causal relation with the occurrence of very large earthquakes. A unified understanding of these slow and fast earthquake processes requires an approach integrating geophysics, seismology, geodesy, geology, and non-equilibrium statistical physics. We welcome presentations based on, but not limited to, geophysical observation, data analysis, analytical theory, numerical simulation, field study, and laboratory experiments.

[SCG53-P27]Nucleation Process of the 2011 *M*_w 6.2 Northern Nagano earthquake

Kengo Shimojo¹, Bogdan Enescu², *Yuji Yagi³, Tetsuya Takeda⁴ (1.Graduate School of Life and Environmental Sciences University of Tsukuba, 2.Graduate School of Science, Kyoto University, 3.Faculty of Life and Environmental Sciences University of Tsukuba, 4.National Research Institute for Earth Science and Disaster Prevention (NIED))

Keywords:Mw 6.2 Northern Nagano earthquake, nucleation process, Matched-Filter Technique, dynamic stress changes, fluid flow, aseismic slip

Introduction. Previous research.

An $M_w6.2$ inland earthquake occurred in northern Nagano region, central Japan, about 13 hours after the $M_w9.0$ Tohoku-oki megathrust earthquake. The regional seismic activity recorded in the Japan Meteorological Agency (JMA) catalog in the first hours following the megathrust event was highly incomplete, thus not allowing a detailed analysis of triggering mechanisms. By applying a Matched Filter Technique (MFT) to the continuous Hi-net (NIED) waveform data, Shimojo et al. (2014) revealed an immediate post-Tohoku seismicity activation in an area located about 10 km south of the $M_w6.2$ Northern Nagano source region. They also detected a few foreshocks close to the hypocenter of the $M_w 6.2$ mainshock, within one hour before the occurrence of the moderate-size event. However, the physical processes that led to the occurrence of the $M_w6.2$ earthquake remained unclear. In this study we take advantage of the data recorded by a dense temporary seismic network operated by NIED from 2008 to 2012 to reveal with an unprecedented resolution the nucleation process that culminated with the occurrence of the Northern Nagano earthquake.

Data and Method

We use the waveform data of the NIED "Hizumi" temporary network, with station spacing of about 5 km or less in the study area. The data recorded by the permanent Hi-net stations (spacing of about 20 km) complements that of the dense regional network. We have first picked P- and S-wave arrivals of earthquakes on the continuous seismograms and use the pick data to locate the events. The earthquakes were then relocated using the tomoDD software (Zhang and Thurber, 2003). The newly

located earthquakes were further used as MFT templates to search for new events within the 13-hour time interval, in the hypocentral region of the M_w 6.2 earthquake.

Results and Discussion

We have detected a total of 285 earthquakes in the source region of the M_w 6.2 event. The earthquakes are relatively small, with magnitudes less than 2.5, and distribute within two spatially distinct clusters: one of these clusters was located close to the hypocenter of the M_w 6.2 event (western cluster), the other about 5 km to the east (eastern cluster).

In the eastern cluster the seismicity starts within one hour after the Tohoku-oki earthquake. The events occur off the M_w 6.2 fault and expand with time from shallow towards deep locations. In the western cluster the seismicity starts immediately after the passage of surface waves excited by a moderate earthquake in the Tohoku-oki aftershock area, which occurred 21 minutes after the M_w 9.0 megathrust; most of these events distribute along the fault line of the Mw6.2 mainshock. Within the western seismicity area, a tight sub-cluster that occurred in the immediate vicinity (within about 2 km) of the M_w 6.2 hypocenter was strongly activated thrice: about 9 hours before, 3 hours before and 50 minutes before the mainshock, respectively. In particular, about 9-hours before and 3-hours before the Nagano earthquake, micro-seismicity migrated towards its epicenter, and just about 1-hour before, microseismicity accelerated within about several hundred meters of the M_w 6.2 epicenter. Migration speeds indicate potential underlying slow-slip culminating with the occurrence of the large inland earthquake, but fluids might have also played a triggering role at a broader, regional scale.

In both western and eastern cluster some events occured immediately after the arrival of surface waves from some $M_{JMA} \ge 5.5$ Tohoku-oki aftershocks. We show that the local seismicity continued intermittently until the occurrence of the M6.2 event, being likely 'modulated' by the arrival of surface waves from larger teleseismic aftershocks off-shore Tohoku.