

---

 [EE] Evening Poster | S (Solid Earth Sciences) | S-CG Complex & General

## [S-CG54]Hard-Rock Drilling: Oman to Oceanic Lithosphere to Island Arc Formation and Beyond

convener: Eiichi TAKAZAWA (Department of Geology, Faculty of Science, Niigata University), Katsuyoshi Michibayashi (Department of Earth and Planetary Sciences, Nagoya University), Peter B Kelemen (共同), Damon A H Teagle (Ocean & Earth Science, National Oceanography Centre Southampton, University of Southampton, SO14-3ZH, Southampton, UK)

Mon. May 21, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

The on-going Oman Drilling Project (OmDP) has drilled numerous hard-rock cores of ancient oceanic lithosphere and the underlying subduction zone of the Samail ophiolite in Oman, with support from ICDP, IODP, the Sloan Foundation's Deep Carbon Observatory, and Japanese, US, and European research agencies. Moreover, a number of IODP expeditions have focused on hard-rock drilling over the last 5 years including Expedition 352 in 2014, which drilled the volcanic sequence associated with subduction initiation in the Bonin fore-arc, Expedition 357, which drilled the Lost City hydrothermal field, Expedition 360, which drilled the lithosphere associated with ultraslow-spreading at the Southwest Indian Ridge in 2016, and Expedition 366, which drilled the serpentinite seamounts in the Mariana fore-arc. In this session, we invite presentations on the scientific results of hard-rock drilling at these and other sites. We also invite related presentations on oceanic lithosphere, island arc formation, and any other significant issue that could be addressed by future hard-rock drilling. This includes marine studies of oceanic lithosphere and on-land geological investigations of ophiolites, accreted arcs, and subduction complexes. The session is intended to be interdisciplinary, including the fields of geophysics, geochemistry, petrology, engineering, and biology.

---

## [SCG54-P06] Overview of Hole GT3A: The sheeted dike/gabbro transition

\*Natsue Abe<sup>1</sup>, Michelle Harris<sup>2</sup>, Katsuyoshi Michibayashi<sup>3</sup>, Juan Carlos de Obeso<sup>4</sup>, Peter B Kelemen<sup>5</sup>, Eiichi TAKAZAWA<sup>6</sup>, Damon A H Teagle<sup>7</sup>, Jude Ann Coggon<sup>7</sup>, Juerg Michael Matter<sup>4</sup>, Phase I Science Party The Oman Drilling Project<sup>8</sup> (1.R&D Center for Ocean Drilling Science Japan Agency for Marine-Earth Science and Technology, 2.Plymouth University, 3.Shizuoka University, 4.Lamont -Doherty Earth Observatory, 5.Columbia University, 6.Niigata University, 7.University of Southampton, Ocean & Earth Science, National Oceanography Centre Southampton, 8.University of Southampton, Oman Drilling Project Office) Keywords: ICDP Oman Drilling Project, sheeted dike/gabbro transition, D/V Chikyu, oceanic crust

Hole GT3A (23.11409 N, 58.21172 E) was drilled by the Oman Drilling Project (OmDP) into Wadi Abdah of the Samail ophiolite, Oman and samples the sheeted dike-gabbro transition zone. OmDP is an international collaboration supported by the International Continental Scientific Drilling Program, the Deep Carbon Observatory, NSF, IODP, JAMSTEC, and the European, Japanese, German and Swiss Science Foundations, with in-kind support in Oman from the Ministry of Regional Municipalities and Water Resources, Public Authority of Mining, Sultan Qaboos University, and the German University of Technology. Hole GT3A was diamond cored in February to March 2017 to a total depth of 400 m. The outer surfaces of the cores were imaged and described on site before being curated, boxed and shipped to the IODP drill ship Chikyu, where they underwent comprehensive visual and instrumental analysis.

Hole GT3A recovered predominantly sheeted dikes and gabbros and has been sub-divided into 4 igneous groups based on the abundance of gabbro downhole. Group 1 (Upper Sheeted Diabase Sequence), group II

(Upper Gabbro Sequence), group III (Lower Sheeted Dike Sequence) and group IV (Lower Gabbro Sequence). Group II and IV are both associated with almost equal proportions of dikes to gabbroic lithologies, whereas group I & III have >95% dikes.

The sheeted dikes were logged as either basalt (46.9 %) or diabase (26.2 %) depending on the predominant grain size of the dike. Gabbroic lithologies include (most to least abundant) gabbro, oxide gabbro and olivine gabbro. Other lithologies present include diorite (7.5%) and tonalite and trondhjemite (1%). Tonalite and trondhjemite are present as cm-sized dikelets and are found within group II and IV. Gabbroic lithologies generally display a varitextured appearance and are characterised by the co-existence of poikilitic and granular domains. Detailed observations of chilled margins and igneous contacts reveal the relative timing of dike and gabbro intrusion, and identify that the Upper Gabbro Sequence intrudes into dikes, whereas the Lower Gabbro Sequence is intruded by dikes. Hydrothermal alteration is recorded throughout the hole and the mean alteration intensity is 54% (range = 10-100%) and mean vein density is 27 veins/m. The intensity of brittle deformation is generally low downhole with restricted intervals associated with fault zones.