

[EE] Evening Poster | S (Solid Earth Sciences) | S-CG Complex & General

[S-CG54]Hard-Rock Drilling: Oman to Oceanic Lithosphere to Island Arc Formation and Beyond

convener: Eiichi TAKAZAWA (Department of Geology, Faculty of Science, Niigata University), Katsuyoshi Michibayashi (Department of Earth and Planetary Sciences, Nagoya University), Peter B Kelemen (共同), Damon A H Teagle (Ocean & Earth Science, National Oceanography Centre Southampton, University of Southampton, SO14-3ZH, Southampton, UK)

Mon. May 21, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

The on-going Oman Drilling Project (OmDP) has drilled numerous hard-rock cores of ancient oceanic lithosphere and the underlying subduction zone of the Samail ophiolite in Oman, with support from ICDP, IODP, the Sloan Foundation's Deep Carbon Observatory, and Japanese, US, and European research agencies. Moreover, a number of IODP expeditions have focused on hard-rock drilling over the last 5 years including Expedition 352 in 2014, which drilled the volcanic sequence associated with subduction initiation in the Bonin fore-arc, Expedition 357, which drilled the Lost City hydrothermal field, Expedition 360, which drilled the lithosphere associated with ultraslow-spreading at the Southwest Indian Ridge in 2016, and Expedition 366, which drilled the serpentinite seamounts in the Mariana fore-arc. In this session, we invite presentations on the scientific results of hard-rock drilling at these and other sites. We also invite related presentations on oceanic lithosphere, island arc formation, and any other significant issue that could be addressed by future hard-rock drilling. This includes marine studies of oceanic lithosphere and on-land geological investigations of ophiolites, accreted arcs, and subduction complexes. The session is intended to be interdisciplinary, including the fields of geophysics, geochemistry, petrology, engineering, and biology.

[SCG54-P11] A new discovery of chromite-hosted inclusions from chromitites of the Samail ophiolite of Oman

*Yuan YAO¹, Eiichi TAKAZAWA^{2,3} (1. Graduate School of Science and Technology, Niigata University, 2. Department of Geology, Faculty of Science, Niigata University, 3. ODS, JAMSTEC)

Keywords: chromitite, inclusions, Oman ophiolite

Podiform chromitites have provided us valuable information on various mantle processes, including melt-mantle reaction, deep-seated magmatic evolution, and mantle dynamics. Varied podiform chromitites have been researched in Oman ophiolite, including the Maqсад chromitite (Arai and Miura, 2016). Many kinds of inclusions have been found in podiform chromitite, Oman ophiolite, such as platinum group element minerals, silicate minerals, even ultra-high pressure minerals, moissanite (Ahmed and Arai, 2003; Trumbull and Yang et al., 2009). The main silicate minerals are hydrous minerals, such as Na-Cr pargasite, aspidolite and so on (Borisova and Ceuleneer et al., 2012; Rospab & Ceuleneer et al., 2017). We studied ore samples from a chromitite mine in the north of Maqсад. The ore samples are massive and banded, mainly consist of disseminated subhedral olivine (0.3~0.5mm) and chromite (0.3~0.5mm), grossular and sulfides are uncommon. Ores are less than 5% alteration, however, some serpentine and native copper and copper carbonate, oxide have been observed in fracture of wall rock dunite. The chemical compositions of chromites range narrow, Cr# (=Cr/[Cr+Al] atomic ratio) is from 0.50 to 0.52, Mg# (=Mg/[Mg+Fe²⁺] atomic ratio) is from 0.57 to 0.64, TiO₂ is from 0.3% to 0.5% which average is 0.4%, FeO is from 17.0% to 22.1%. Different types of inclusions were found in chromites. The silicate inclusions range in diameter from 5 μm to 200 μm, contain pargasite, aspidolite, high Cr# (Cr# > 60) chromite, diopside, enstatite, forsterite, grossular, anorthite (An=99), titanite, pentlandite,

chlorite. The chemical compositions of chromites may represent the melt composition from which they formed, the inferred parental melt composition in equilibrium with the chromitites has high Al_2O_3 content (14.9–15.4 wt.%), which is very similar to MORB magmas (Al_2O_3 ~15wt.%; Kamenetsky et al., 2001). Considering the magmatic anorthite is a typical mineral of gabbro in the arc setting (Beard, 1986), it may indicate the chromite record MORB-like and arc-like melts. Moreover, the hydrous silicate inclusions homogenized experiment has been carried out, at 900°C, 1100°C, 1150°C, 5mm cube samples are heated by muffle furnace, keep the temperature for 2 min then quenching by water; at 1300°C, chromite grains are heated by thermal gravimetric analyzer, cooling by air. Homogenized volume of inclusions increased with temperature such as 5% (2 homogenized inclusions), 10% (5 homogenized inclusions), 50% (only pargasite, diopside still exist), 100% from 900°C to 1300°C, respectively. Moreover, the compositions of homogenized inclusions are various, the contents of SiO_2 , FeO and TiO_2 range from 14% to 71%, from 0.6% to 40.4%, from 0.2% to 6%, respectively, which indicates the heterogeneity of the silicate melt when they were trapped by chromite. It can be inferred that more than one simple melt interacted with the mantle sequences. Further investigation is required to prove this hypothesis.

Ahmed, A.H. and Arai, S., 2003. *The Canadian Mineralogist*, 41(3): 597-616.

Arai, S. and Miura, M., 2016. *Lithos*, 264: 277-295.

Beard, J.S., 1986. *Geology*, 14(10): 848-851.

Borisova, A.Y. et al., 2012. *Journal of Petrology*, 53(12): 2411-2440.

Rospabé, M., Ceuleneer, G., Benoit, M., Abily, B. and Pinet, P., 2017. *Geology*, 45(5): 471-474.

Kamenetsky, V.S., Crawford, A.J., Meffre, S., 2001. *Journal of Petrology* 42, 655–671.

Trumbull, R.B. et al., 2009. *Lithos*, 113(3-4): 612-620.