
[EJ] Evening Poster | S (Solid Earth Sciences) | S-CG Complex & General

[S-CG57]Dynamics in mobile belts

convener:Yukitoshi Fukahata(Disaster Prevention Research Institute, Kyoto University), Toru Takeshita(Department of Natural History Sciences, Graduate School of Science, Hokkaido University), Hikaru Iwamori(海洋研究開発機構・地球内部物質循環研究分野)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

The dynamic behaviours of mobile belts are expressed across a wide range of time scales, from the seismic and volcanic events that impact society during our lifetimes, to orogeny and the formation of large-scale fault systems which can take place over millions of years. Deformation occurs on length scales from microscopic fracture and flow to macroscopic deformation to plate-scale tectonics. To gain a physical understanding of the dynamics of mobile belts, we must determine the relationships between deformation and the driving stresses due to plate motion and other causes, which are connected through the rheological properties of the materials. To understand the full physical system, an integration of geophysics, geomorphology, and geology is necessary, as is the integration of observational, theoretical and experimental approaches. In addition, because rheological properties are greatly affected by fluids in the crust and fluid chemical reactions, petrological and geochemical approaches are also important. After the 2011 great Tohoku-oki earthquake, large-scale changes in seismic activity and regional scale crustal deformation were observed, making present-day Japan a unique natural laboratory for the study of the dynamics of mobile belts. This session welcomes presentations from different disciplines, such as seismology, geodesy, tectonic geomorphology, structural geology, petrology, and geofluids, as well as interdisciplinary studies, that relate to the dynamic behaviour of mobile belts.

[SCG57-P12]Mechanism of vertical displacement on the Northeast Japan fore-arc region through gigantic earthquake cycles

*Ryohei Sasajima¹, Bunichiro Shibazaki¹, Hikaru Iwamori^{2,3}, Takuya Nishimura⁴, Yoshihiko Nakai¹

(1.International Institute of Seismology and Earthquake Engineering, Building Research Institute, 2.Japan Agency for Marine-Earth Science and Technology, 3.School of Science, Tokyo Institute of Technology, 4.Disaster Prevention Research Institute, Kyoto University)

Keywords:Vertical deformation through gigantic earthquake cycles, Vertical displacement paradox, 2011 Tohoku earthquake, Kinematic modeling of crustal deformation, Finite element method

Fore-arc region in the middle (38-39.5°N) Northeast Japan (NEJP) exhibited fast subsidence (up to 3-4 mm/yr) in the interseismic period during ~100 years before the 2011 Tohoku earthquake (M_w 9) (e.g., Nishimura, 2014). At the 2011 Tohoku earthquake, the fore-arc region also subsided by 0.4-1.2 m (Nishimura, 2014). On the other hand, long-term vertical displacement rate on the fore-arc region indicates very slow uplift (0.1-0.2 mm/yr) (Tajikara, 2004). Therefore, when and how the vertical displacement on the fore-arc region balances and satisfies the long-term displacement through the gigantic earthquake cycle has been argued as the vertical displacement paradox and is still controversial (e.g., Ikeda, 2014; Nishimura, 2014; Sagiya, 2015; Hashima and Sato, 2017). This study attempted to elucidate the mechanism of the vertical displacement on the NEJP fore-arc region through the gigantic earthquake cycle and solve the vertical displacement paradox via a numerical modeling.

This study modeled the crustal deformation through the gigantic earthquake cycles using finite element method (Shibazaki *et al.*, 2007) considering heterogeneous rheological structure. The interplate coupling

and the earthquake slip on the asperity of the gigantic earthquake were modeled via the slip-node method (Melosh and Raefsky, 1981) based on the back-slip model (Savage, 1983). We modeled the fault creeping on the middle and deeper portions of the plate interface via a thin low-viscous layer (e.g., Hu *et al.*, 2016). The model considered a ductile shear zone, which is extended from the deeper portion of the plate interface and connects to the viscoelastic mantle wedge.

The modeling result indicated that interseismic subsidence rate in the fore-arc region increased with the duration of locking of the asperity of the 2011 Tohoku earthquake. It was caused by increasing slip deficit rate on the deeper portion of the plate interface caused by continuous locking of the asperity during several hundreds years with viscoelastic relaxation in the mantle wedge and the ductile shear zone. Therefore, the result implies that the observed fast subsidence during ~100 years before the 2011 Tohoku earthquake might occur only in the later period of the gigantic earthquake cycle.

The modeling result indicated that large postseismic uplift occurred on the fore-arc region during ~50 years after the gigantic earthquake caused by large afterslip (up to ~10 m) with long duration, which was controlled by postseismic viscoelastic shear deformation in the ductile shear zone. The result predicts that the coseismic fore-arc subsidence will almost recover 40-50 years after the 2011 Tohoku earthquake mainly due to this long duration afterslip. The sum of the large postseismic uplift (~0.4-1.4 m/50 yr) caused by the long-duration afterslip and subsequent uplift in the earlier interseismic period (~0.6-0.8 m/150 yr) will balance to the sum of the fast subsidence in the later period of the gigantic earthquake cycle (~0.6-0.8 m/300 yr) and the coseismic subsidence (~0.4-1.4 m). This is the possible mechanism to solve the vertical displacement paradox in the NEJP fore-arc region.

Similar fast interseismic subsidence also has been observed in the East Hokkaido fore-arc region (western Kuril subduction zone), where the gigantic earthquake occurred in the 17th century (e.g., Nanayama *et al.*, 2003) and subsequent large postseismic uplift with long-duration (~1.5 m/60 yr) was estimated (Sawai *et al.*, 2004). Our result implies that this fast subsidence may also be caused by the continuous locking of the asperities of the gigantic earthquake during several hundreds years, i.e., at least ~30 m of the slip deficit has been cumulated on the asperities of the 17th century event.