[EJ] Evening Poster | S (Solid Earth Sciences) | S-CG Complex & General

[S-CG59]Structure and evolution of Japanese islands - Formation of island arc systems and earthquake cycles

convener:Hiroshi Sato(Earthquake Prediction Research Center, Earthquake Research Institute, The University of Tokyo), Masanao Shinohara(Earthquake Research Institute, University of Tokyo), Masahiro Ishikawa(横浜国立大学大学院環境情報研究院, 共同), Makoto MATSUBARA(National Research Institute for Earth Science and Disaster Resilience)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe) Subduction processes such as accretion, back-arc-spreading, and arc-arc collisions have shaped the Japanese island arc. Recent advances in seismic imaging, both passive and controlled source, have produced new images of the crust-mantle structure under Japan and surrounding regions. Through the influence of pre-existing faults and rheological structures, these crust and mantle structures are exerting strong control on active tectonic processes like seismic activity and crustal deformation in the overriding plate. We seek contributions that document and/or model the deformation of the Japanese islands over a variety of time scales from the earthquake cycle to the tectonic evolution of the Japanese island arc, and from a range of research fields including seismology, geology, geochemistry, tectonic geomorphology, and geodynamics. Multidisciplinary studies are encouraged. We also welcome contributions in numerical or analogue geodynamical modeling that explore deformation processes.

[SCG59-P01]Scattered wavefield imaging of the Japan subduction structure using Hi-net array data

*Younghee Kim¹, Hitoshi Kawakatsu² (1.Seoul National University, 2.Earthquake Research Institute, University of Tokyo)

Keywords: Teleseismic migration, Subduction-zone geometry, Hi-net data

The Japan arcs exhibit significant along-arc transition in slab geometry, subduction-driven magmatic activity, and plate coupling state. We exploit teleseismic data from dense Hi-net array to define the present-day subduction-zone geometry and to constrain the plate boundary dynamics in response to variable slab geometries and physical properties of the subducting material. We retrieve backscattered (or reverberated) body-wave phases in addition to direct Ps (incident P wave forward-scattered as an S wave) and apply the teleseismic migration method based on the Generalized Radon Transform (Bostock et al., 2001) to accurately define both Pacific and Philippine Sea plate subduction structures. This method inverts the scattered waves for sharp variations in the Earth's elastic properties beneath the high-density array using analytical expressions for their travel-times and amplitudes (Bostock et al., 2001). The four scattering modes (Ps, PpPs, PpSs|v, and PpSs|h) are sensitive to S-wave velocity perturbations (dVs/Vs) and are combined to form a single composite image. As more migrated phases are stacked in, the artifacts due to cross-mode contamination (i.e., parallel echoes of the real structure) become attenuated while energy mapped to its correct depth is sharply imaged. Preliminary images show complex slab structures down to a depth of 200 km with significant along-arc velocity variations at the top of the plate. Such differences in the velocity structures may indicate different hydration state due to slab age and/or presence of low-strength materials.