
[JJ] Evening Poster | S (Solid Earth Sciences) | S-EM Earth's Electromagnetism

[S-EM17]Geomagnetism, Paleomagnetism and Rock Magnetism

convener:Nobutatsu Mochizuki(Priority Organization for Innovation and Excellence, Kumamoto University),
Hisayoshi Shimizu(Earthquake Research Institute, University of Tokyo)

Mon. May 21, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

We are going to discuss the issues on the magnetic fields of the Earth and planets, paleomagnetism, rock-magnetism, and their applications. This session includes the following topics: (1) observation and analysis of the magnetic fields of the Earth and planets, (2) paleomagnetic field variations obtained from natural and archaeological materials, (3) numerical simulations on the magnetic fields of the Earth and planets, (4) measurements and theories of magnetic properties of rocks, minerals, meteorites and other materials, (5) climate changes and global and local surface tectonics based on the paleomagnetic measurements of rocks and sediments, (6) observations of the magnetic anomalies and the crustal magnetization models of the Earth, planets and satellites, and (7) developments of the experimental method and data analysis. The presentation and discussion will be made in Japanese or English in this session.

[SEM17-P02]Spatial distribution of regularity of local geomagnetic jerks and electrical conductivity at the bottom of the mantle

*Hisayoshi Shimizu¹, Hisashi Utada¹ (1.Earthquake Research Institute, University of Tokyo)

Keywords:geomagnetic jerk, electrical conductivity

Geomagnetic jerk, abrupt change of geomagnetic secular acceleration, is considered to have the shortest time-scale among the observed geomagnetic main field variations. Geomagnetic jerks are observed around 1969, 1978, and some other years, and they have been analyzed to estimate or to discuss the electrical conductivity in the mantle (e.g. Backus, 1983; Alexandrescu et al., 1999; Nagao et al., 2003; Pinheiro and Jackson, 2008). For example, Alexandrescu et al. (1999) applied wavelet analysis to the observed magnetic field components to identify geomagnetic jerks and their regularity. The observed regularity at geomagnetic stations were used to estimate the regularity of magnetic field variation at the core-mantle boundary and the electrical conductivity of the lower mantle.

Geomagnetic jerks were observed around 2003 and 2007 but the signature of them are strong only in the south Atlantic region (e.g. Chulliat et al., 2010). We showed in a previous work that the regularity of the local geomagnetic jerk around 2007 at Mbour (MBO, Senegal) is significantly higher than that of the global geomagnetic jerks, and suggested that the local geomagnetic jerk can be generated due to the combined effect of toroidal magnetic field variation at the CMB and the electrical heterogeneity at the bottom of the mantle. In this presentation, we show the spatial distribution of the regularity of local geomagnetic jerks and discuss electrical conductivity heterogeneity in the deep mantle.

Reference:

- Alexsandrescu, M. et al., 1999, JGR, 104, 17,735-17,745.
- Backus, G., 1983, GJRastS, 74, 713-746.
- Chulliat, A. et al., 2010, GRL, 37, L0730.
- Nagao, H. et al., 2003, JGR, 108, 2254.
- Pinheiro and Jackson, 2008, GJI, 173, 781-792.