
[EE] Evening Poster | S (Solid Earth Sciences) | S-IT Science of the Earth's Interior & Tectonophysics

[S-IT18]Planetary cores: Structure, formation, and evolution

convener: Hidenori Terasaki (Graduate School of Science, Osaka University), Eiji Ohtani (Department of Earth and Planetary Materials Science, Graduate School of Science, Tohoku University), William F McDonough (共同), George Helffrich (Earth-Life Science Institute, Tokyo Institute of Technology)

Mon. May 21, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

There are fundamental links between the formation and evolution of planets and their satellites to that of their cores. Defining the physical and chemical properties of the cores of these terrestrial bodies are fundamental for understanding their internal structures and thermal profile. Recent advances in experimental and theoretical studies provide new insights and applications to the Earth's cores and other terrestrial bodies. Future exploration missions will obtain data on the internal structure of terrestrial planets (e.g., Mars and Mercury) and planet-satellite systems. We anticipate presentations on recent advances on the physical and chemical properties of cores and discussions regarding the latest views of their formation and evolution. We welcome contributions from mineral/rock physics, geophysics, geochemistry, geodynamics, and planetary science.

[SIT18-P05]Stratification of the Earth's core by SiO₂ crystallization

*George Helffrich¹, Guillaume Morard³, Kei Hirose^{1,2}, Satoshi Kaneshima⁴ (1.Earth-Life Science Institute, Tokyo Institute of Technology, 2.Department of Earth and Planetary Science, University of Tokyo, 3.IMPMC, Universite Pierre et Marie Curie - Paris 6, 4.Department of Earth and Planetary Sciences, University of Kyushu)

Keywords: core, stratification, secular variation, seismic wavespeeds

The topmost ~300 km of the Earth's core features slightly lower wavespeeds than the one dimensional radial earth model PREM. One way this could arise is if this lower speed layer represents a crystallization boundary inside the core, similar to the cloud base in the atmosphere where water-saturated air condenses out water. In the case of the Earth's core, saturation in Si+O could crystallize SiO₂ that rises upward to the CMB and leaves the core due to SiO₂ being even less dense than the base of the Earth's present mantle. This mechanism addresses a criticism of stable core stratification arising from study of the secular variation of the Earth's magnetic field, which seems to require radial fluid motions that strict stratification forbids. Motion of core liquid through a saturation front, which changes its composition, temperature and wavespeeds, facilitates patterns of secular variation attributable to radial motion. We explore this scenario by finding SiO₂ saturation parameters consistent with experimental partitioning of Si and O in metal, and derive the temperature and core compositions compatible with the observed wavespeed profile in the outer core.