

[EE] Evening Poster | S (Solid Earth Sciences) | S-IT Science of the Earth's Interior & Tectonophysics

## [S-IT20] Structure and Dynamics of Earth and Planetary Mantles

convener: Takashi Yoshino (Institute for Planetary Materials, Okayama University), Dapeng Zhao (Department of Geophysics, Tohoku University), Takashi Nakagawa (海洋研究開発機構数理科学・先端技術研究分野)

Mon. May 21, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Interdisciplinary approach can lead to a better understanding of dynamics and evolution of the deep interiors of the Earth and planets. We welcome any submissions of recent results in observational, theoretical and experimental studies on seismology, geomagnetism, mineral physics, dynamics of deep interiors, and any other relevant fields from researchers in many countries. Integration of such results is also welcome. In particular, we encourage any contributions focusing on "plate and mantle dynamics in Earth and terrestrial planets".

## [SIT20-P06] Effects of pressure and water on electrical conductivity of carbonate melt with implications for conductivity anomaly in continental mantle lithosphere

\*Takashi Yoshino<sup>1</sup>, Benjamin Gruber<sup>2</sup>, Clayton Reinier<sup>3</sup> (1. Institute for Planetary Materials, Okayama University, 2. Earth and Atmospheric Sciences, University of Alberta, 3. Department of Geology and Geography, University of West Virginia)

Keywords: Electrical conductivity, Pressure

Electrical conductivity of Na, Mg-bearing carbonate melts was measured in a Kawai-type multi-anvil apparatus as a function of pressure. The carbonate samples were mixtures of  $\text{MgCO}_3$  and  $\text{Na}_2\text{CO}_3$  or  $\text{Mg}_5(\text{CO}_3)_4(\text{OH})_2 \cdot 4(\text{H}_2\text{O})$  and  $\text{Na}_2\text{CO}_3$ . High pressure experiments on the carbonate systems were performed up to 1800 K in a wide pressure range from 3.4 to 10.9 GPa. The sample conductivity abruptly changes at eutectic temperature, which increases with increasing pressure. Hydrous carbonate yields lower eutectic temperature than anhydrous carbonate and has weaker pressure dependence. Molten state carbonates show a very high electrical conductivity with temperature dependence following an Arrhenius law. As pressure increases, the conductivity decreases. The negative pressure dependence on electrical conductivity of hydrous carbonate melt is larger than that of the anhydrous one. Activation volumes were determined to be  $\Delta V = 1.81$  and  $3.61 \text{ cm}^3 \cdot \text{mol}^{-1}$  for anhydrous and hydrous carbonate melts, respectively. The high electrical conductivity observed in the mantle beneath the Slave and Brazilian cratons can be explained by the process of lithospheric rejuvenation due to small amounts of hydrous carbonated melt released from crystallization of kimberlitic magma at the base of continental mantle lithosphere.