

---

 [EE] Evening Poster | S (Solid Earth Sciences) | S-IT Science of the Earth's Interior & Tectonophysics

## [S-IT22]Interaction and Coevolution of the Core and Mantle in the Earth and Planets

convener: Tsuyoshi Iizuka (University of Tokyo), Hidetoshi Shibuya (Department of Earth and Environmental Sciences, Faculty of Advanced Science and Technology, Kumamoto University), Taku Tsuchiya (愛媛大学地球深部ダイナミクス研究センター, 共同), Kenji Ohta (Department of Earth and Planetary Sciences, Tokyo Institute of Technology)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Recent observational and experimental investigations have significantly advanced our understanding of the structure and constituent materials of the deep Earth. Yet, even fundamental properties intimately linked with formation and evolution of the planet, such as details of the chemical heterogeneity in the mantle and light elements dissolved in the core, remained unclear. Seismological evidence has suggested a vigorous convection in the lower mantle, whereas geochemistry has suggested the presence of stable regions there that hold ancient chemical signatures. The amounts of radioactive isotopes that act as heat sources and drive dynamic behaviors of the deep Earth are also still largely unknown. We provide an opportunity to exchange the achievements and ideas, and encourage persons who try to elucidate these unsolved issues of the core-mantle evolution using various methods, including high-pressure and high-temperature experiments, high-precision geochemical and paleomagnetic analyses, high-resolution geophysical observations, geo-neutrino observations, and large-scale numerical simulations. Since this session is co-sponsored by geomagnetism, paleomagnetism and rock magnetism division of the SGPSS, contributions in geomagnetism and geodynamo simulation are also encouraged.

---

### [SIT22-P07]Lattice thermal conductivity of (Mg,Fe)O

\*Akira Hasegawa<sup>1</sup>, Kenji Ohta<sup>1</sup>, Takashi Yagi<sup>2</sup>, Kei Hirose<sup>1</sup>, Tadashi Kondo<sup>3</sup> (1.Tokyo Institute of Technology, 2.National Institute of Advanced Industrial Science and Technology, 3.Osaka University)

Keywords: thermal conductivity, lower mantle, magnesio-wüstite, wüstite

The Earth has been cooling since it was born about 4.6 billion years ago. To decipher the thermal history of the Earth, thermophysical properties of the lower mantle materials that constitute more than half the volume of the Earth is of great importance. A number of research has suggested the possibility that (Mg,Fe)O ferropericlase, one of the main constituent minerals of the lower mantle, exists with very iron-rich chemical composition at the Earth's core-mantle boundary (CMB). Such iron-rich (Mg,Fe)O magnesio-wüstite at the CMB may cause regional variation of thermal conductivity of the lowermost mantle due to its distinct iron concentration, which potentially influences the mantle convection style, inner core age, inner core structure, geomagnetic field reversal frequency and so on [e.g. Olson, 2016]. However, there is no systematic study to examine the effect of iron on the thermal conductivity of (Mg,Fe)O solid solution under high pressure. In this study, we measured lattice thermal conductivity of FeO wüstite and (Mg,Fe)O magnesio-wüstite with various iron contents at high pressures, and evaluated their pressure and compositional dependence. As a results, the thermal conductivity of FeO decreased in the range of 33 to 60 GPa where is in a stability field of the rB1 structure. In the B1 and rB1 structures, the pressure dependence of the thermal conductivity is obviously different. In order to evaluate the effect of iron incorporation on thermal conductivity, we measured lattice thermal conductivity of (Mg,Fe)O which has same crystal structure as CMB conditions. Our results show much lower lattice conductivity of iron-rich magnesio-wüstite than that of MgO and FeO due to strong iron impurity phonon scattering, which would help to estimate the thermal

conductivity of the expected iron-rich region in the lowermost mantle.

Reference: Olson, P. Mantle control of the geodynamo: Consequences of top-down regulation, *Geochem. Geophys. Geosys.* 17, 1935–1956, (2016).