## [EE] Evening Poster | S (Solid Earth Sciences) | S-IT Science of the Earth's Interior & Tectonophysics

## [S-IT22]Interaction and Coevolution of the Core and Mantle in the Earth and Planets

convener:Tsuyoshi lizuka(University of Tokyo), Hidetoshi Shibuya(Department of Earth and Environmental Sciences, Faculty of Advanced Science and Technology, Kumamoto University), Taku Tsuchiya(愛媛大学地球深部ダイナミクス研究センター, 共同), Kenji Ohta(Department of Earth and Planetary Sciences, Tokyo Institute of Technology)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe) Recent observational and experimental investigations have significantly advanced our understanding of the structure and constituent materials of the deep Earth. Yet, even fundamental properties intimately linked with formation and evolution of the planet, such as details of the chemical heterogeneity in the mantle and light elements dissolved in the core, remained unclear. Seismological evidence has suggested a vigorous convection in the lower mantle, whereas geochemistry has suggested the presence of stable regions there that hold ancient chemical signatures. The amounts of radioactive isotopes that act as heat sources and drive dynamic behaviors of the deep Earth are also still largely unknown. We provide an opportunity to exchange the achievements and ideas, and encourage persons who try to elucidate these unsolved issues of the core-mantle evolution using various methods, including high-pressure and hightemperature experiments, high-precision geochemical and paleomagnetic analyses, high-resolution geophysical observations, geo-neutrino observations, and large-scale numerical simulations. Since this session is co-sponsored by geomagnetism, paleomagnetism and rock magnetism division of the SGEPSS, contributions in geomagnetism and geodynamo simulation are also encouraged.

## [SIT22-P32]Study of the required Rayleigh number to sustain geodynamo with various inner core radius

\*Nishida Yuki<sup>1</sup>, Yuto Katoh<sup>1</sup>, Hiroaki Matsui<sup>2</sup>, Masaki Matsushima<sup>3</sup>, Atsushi Kumamoto<sup>1</sup> (1.Department of Geophysics, Graduate School of Science, Tohoku University, 2.Dept. of Earth and Planetary Sciences, University of California, Davis, 3.Department of Earth and Planetary Sciences, School of Science, Tokyo Institute of Technology)

Keywords:geodynamo, the past Earth, the inner core size

The Earth has an intrinsic magnetic field. Paleomagnetic studies suggest that the geomagnetic field has been sustained over 3.5 billion years [e.g., Biggin et al., 2015], and studies of the thermal history of the Earth suggest that the inner core grows to the current size in the last billion years [e.g., O'Rourke &Stevenson, 2016]. The magneto-fluid convection driven by thermal and compositional buoyancy in the outer core is thought to cause geodynamo. While characteristics of magneto-fluid convection have been investigated in detail for the present radius ratio  $r_i/r_o=0.35$ , there have been few simulations for the settings of  $r_i/r_o < 0.35$ , which correspond to the environment of the past Earth. It is important to investigate the dynamo process occurring under the settings of  $r_i/r_o < 0.35$ , so as to examine conditions of the past Earth. Using the numerical dynamo open source code Calypso [Matsui et al., 2014], we perform a series of numerical simulations of the thermal convection and dynamo with the smaller inner core settings than the present inner core. To investigate the characteristics of the convection. Only *Ra* is treated as a variable and the other control parameters, the Ekman number *E*, the Prandtl number *Pr*, and the magnetic Prandtl number *Pm* are fixed to be  $E = 10^{-3}$ , Pr = 1, and Pm = 5, respectively.

In previous dynamo simulations, the property of *Ra* around the onset of dynamo action has been revealed for various radius ratios [Heimpel et al., 2005]. The tendency of the dynamo action has also been investigated by previous studies in the present ratio  $r_i/r_o = 0.35$  [Christensen and Aubert, 2006], but the behavior of dynamo with Ra has not been fully understood with the smaller inner core. We perform dynamo simulations with a range of 1.9  $Ra_{crit} < Ra < 9.7 Ra_{crit}$ , where  $Ra_{crit}$  is the critical Rayleigh number, and with the aspect ratio  $r_i/r_o = 0.15$ , 0.25, and 0.35, to quantitatively understand the tendencies with the smaller inner core. The results with  $r_i/r_o = 0.25$  show that the sustained dipolar dominant dynamo occurs just above the onset of the dynamo action. In the case with larger Ra than that for the dipolar regime, non-dipolar components of the magnetic field is sustained. Finally, dynamo failed in the case with Ra >8.1 Ra<sub>crit</sub>. The simulation results reveal that the magnetic energy density is largest at the dynamo onset, and becomes smaller at larger Ra. This implies that dynamo is not likely to be sustained in intense convection under the assumed setting. The results of  $r_i/r_o = 0.25$  and 0.35 imply that the revealed tendency revealed by the present study can be applied to  $r_i/r_o = 0.15$ . Comparing the simulation results in the same  $Ra/Ra_{crit}$  (= 3.6), the magnetic dipole moment becomes smaller with the smaller inner core. This suggests a possibility that the convection favorable for the dipolar dominant dynamo is difficult to occur with the smaller inner core. We also find that the Ra range of the dipolar dominant cases becomes narrower with the smaller inner core, implying that Ra was very selective at the past Earth. To interpret the simulation results with the real physical quantities, the average temperature difference between the core-mantle boundary and the inner core boundary, ΔT, is estimated. Δ T in  $r_i/r_o = 0.25$  is larger than that in  $r_i/r_o = 0.35$ , so required buoyancy to sustain a dipolar dominant dynamo is larger with the smaller inner core.