

[JJ] Evening Poster | S (Solid Earth Sciences) | S-MP Mineralogy & Petrology

[S-MP38]Physics and Chemistry of Minerals

convener:Hiroaki Ohfuji(Geodynamics Research Center, Ehime University), Seiji Kamada(Frontier Research Institute for Interdisciplinary Sciences, Tohoku University)

Thu. May 24, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

In this session, we will discuss the physics and chemistry of the Earth and planetary materials (including amorphous and melts) based on the results obtained from various experimental methods such as X-ray diffraction, FT-IR, Raman spectroscopy, electron microscopy and computer simulations.

[SMP38-P02]Structure changes of hydromagnesite $Mg_5(CO_3)_4(OH)_2 \cdot 4H_2O$ with temperature and pressure*Gen-ichiro Yamamoto¹, Atsushi Kyono¹, Tomoya Tamura¹, Sota Takagi¹ (1.University of Tsukuba)Keywords:CO₂ geological storage, hydromagnesite $Mg_5(CO_3)_4(OH)_2 \cdot 4H_2O$, structure change, high-temperature XRD, high-pressure XRD

The global warming has become one of the most important environment problems in recent years. Geological storage of CO₂ has great potential to eliminate atmospheric CO₂ emitted by fossil fuel combustion (XUE and Nakao 2008). Regarding mineral CO₂ sequestration, CO₂ is chemically stored through mineral carbonation in solid carbonates. Magnesium and/or calcium carbonate minerals such as magnesite MgCO₃, dolomite CaMg(CO₃)₂, calcite CaCO₃, and aragonite CaCO₃ have long been recognized as the best CO₂ storage. Because hydrous magnesium carbonates can be directly precipitated from aqueous solution in a range between 0 and 200 °C (Cheng et al. 2009; Yamamoto et al. 2017), we consider that hydrous magnesium carbonates would be the best candidate for the CO₂ storage process. The major hydrous magnesium carbonates formed from aqueous solution are nesquehonite MgCO₃·3H₂O and hydromagnesite Mg₅(CO₃)₄(OH)₂·4H₂O. The physical properties of these minerals are critically important factors to assess environmental safety and phase stability over geological time scale, but they have not been fully investigated yet, especially of hydromagnesite. In this study, we performed investigated the structural behaviors of hydromagnesite under high temperature and high pressure using HT-XRD, TG-DTA, HP-XRD to reveal the physical properties and structural stability of hydromagnesite.

The results of *in-situ* high-temperature XRD showed that, no significant change was observed up to 170 °C. With increasing temperature, however, the intensities started to decrease at 200 °C, and all peaks disappeared at 290 °C. Above the decomposition temperature a few peaks corresponding to periclase appeared. The results of TG-DTA clearly showed two weight loss steps in the temperature range from 200 to 340 °C and from 340 to 500 °C, which correspond to the dehydration and decarbonation of hydromagnesite, respectively. Therefore, hydromagnesite decomposes into periclase, carbon dioxide, and water as following reaction: $Mg_5(CO_3)_4(OH)_2 \cdot 4H_2O \rightarrow 5MgO + 4CO_2 + 5H_2O$. No magnesite is formed from hydromagnesite with temperature. The results of *in-situ* high-pressure XRD showed that the intensities gradually decreased with compression, and then most of peaks disappeared above 6.4 GPa. After decompressing to 4.2 GPa, however, the hydromagnsite reappeared. It suggests that no dehydration occurs below at least 6.4 GPa. The amorphousization of hydromagnesite with compression would be caused by disordered arrangements of MgO₆ octahedra and CO₃ triangles in the structure.