

[JJ] Evening Poster | S (Solid Earth Sciences) | S-SS Seismology

## [S-SS12]Seismicity

convener:Kei Katsumata(Institute of Seismology and Volcanology, Hokkaido University)

Thu. May 24, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

This session aims to improve our understanding on seismicity. Any contribution on behavior of earthquakes as a cluster, such as regional seismicity and aftershocks, are welcomed. We also welcome contribution on temporal and spatial interactions that control seismicity, and tectonic processes, and geological and thermal structures that regulate seismicity.

## [SSS12-P01]Ocean bottom seismic observation around erupting

### Nishinoshima and Chichijima, Ogasawara Islands, Japan

\*Noritaka Hanamura<sup>1</sup>, Kenji Nakata<sup>2</sup>, Hisatoshi Baba<sup>1</sup>, Kazuhiro Kimura<sup>3</sup>, Yutaka Nagaoka<sup>2</sup> (1.Tokai University Graduate School of Oceanography Course of Oceanography, 2.Seismology and Tsunami Research Department,Meteorological Research Institute(MRI), 3.Seismology and Volcanology Department,JMA)

Keywords:Nishinoshima, Ogasawara, Volcano, Earthquake

Nishinoshima Island located in Ogasawara Islands started volcanic eruption in November 2013. Meteorological Research Institute carried out ocean bottom seismic observation from July to September 2015 to detect earthquakes associated with the eruption of Nishinoshima, and to improve hypocenter determination accuracy in the area of the Ogasawara Islands. Ocean Bottom Seismographs (OBSs) were installed at five sites around Nishinoshima, and seven sites around Chichijima Island. This study aims to clarify the results of earthquake hypocenter distribution and analyze them.

Around Chichijima area (July 15 to September 30, 2015)

(1) 223 hypocenters were determined, and these were distributed along the plate boundary of Hayes et al.(2012).

(2) These hypocenters were distributed like “double seismic planes (Double Seismic Zone)” in depth ranges of 90-180 km in the subducting Pacific slab.(Nakata et al.,2017)

(3) Magnitudes of determined hypocenters were distributed in M1.8-M5.0, and relatively large (M4.0-M5.0) earthquakes were distributed in west side of the observation network.

Around Nishinoshima area (September 1 to 3, 2015)

(1) 318 hypocenters were determined, and they were distributed at depth shallower than 5.0 km.

(2) Most of the hypocenters were identified at depth between 0 km and 3.0 km. This depth range corresponds to the area just above the magma chamber which Sano et al.(2016) suggested with temperature of 970-990 degrees at the depth of 3.0-6.0 km beneath Nishinoshima Island.

(3) Magnitudes of the determined hypocenters were distributed in M0.3-M1.2, and relatively large earthquakes were located in deeper area.

In addition to the above results, we report spectrum analysis of seismic records in detail.