[JJ] Evening Poster | S (Solid Earth Sciences) | S-SS Seismology

[S-SS12]Seismicity

convener:Kei Katsumata(Institute of Seismology and Volcanology, Hokkaido University) Thu. May 24, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe) This session aims to improve our understanding on seismicity. Any contribution on behavior of earthquakes as a cluster, such as regional seismicity and aftershocks, are welcomed. We also welcome contribution on temporal and spatial interactions that control seismicity, and tectonic processes, and geological and thermal structures that regulate seismicity.

[SSS12-P02]Focal mechanisms of the double seismic zone in the Pacific plate, off Boso: Change before and after the 2011 Tohoku EQ

Tomoharu Sato¹, *Toshinori Sato², Asami Terada³, Masanao Shinohara⁴, Kimihiro Mochizuki⁴, Tomoaki Yamada⁴, Kenji Uehira⁵, Takashi Shimbo⁶, Shuichi Kodaira⁶, Yuya Machida⁶, Ryota Hino⁷, Ryosuke Azuma⁷, Yoshio Murai⁸, Yoshihiro Ito⁹, Hiroshi Yakiwara¹⁰, Kenji Hirata⁵ (1.Faculty of Science, Chiba Univ., 2.Graduate School of Science, Chiba Univ., 3.Kozo Keikaku Engineering, 4.ERI, Univ. Tokyo, 5.NIED, 6.JAMSTEC, 7. Tohoku Univ., 8.Hokkaido Univ., 9.DPRI, Kyoto Univ., 10.Kagoshima Univ.) Keywords:double seismic zone, focal mechanism, the Pacific plate, off the Boso Peninsula, the 2011 Tohoku Earthquake, stress state

1.Introduction

The Pacific plate and the Philippine Sea plate subduct off the Boso Peninsula, where many seismic events such as 1923 Taisho EQ, 1677 Enpo EQ, and slow slip events occurred. It is important to investigate seismic structures and seismicities at this region. Terada et al. (JpGU, 2016) revealed a 3D seismic structure and seismicity using data from land and ocean bottom observations. They showed a double seismic zone in the Pacific plate off the Boso Peninsula. This presentation shows focal mechanisms of the double seismic zone, and discusses stress state of the Pacific plate, especially, before and after the 2011 Tohoku Earthquake.

2. Data and analysis

The used data are waveform data from Hi-net by NIED, 2009-2010 ocean bottom observations by the R/V Hakuho-maru (KH09-3, etc.), and the aftershock survey of the 2011 Tohoku EQ using ocean bottom seismometers (2011/3/15-6/30). We select events in the double seismic zone, and pick polarities of the P wave first motions. The depth of the selected events are about 70-140 km. We calculate emergent azimuth and dip of each event-receiver pair using the 3D structure determined by Terada et al. (JpGU, 2016), and search optimal focal mechanisms using a grid search method.

3. Results

Many focal mechanisms before the 2011 Tohoku EQ have down dip compression for the upper surface events, and down dip extension for the lower surface events. This stress state may represent an unbending of the Pacific plate at the depth. On the other hand, many mechanisms after the 2011 Tohoku EQ have down dip compression for both upper and lower surface events. This may indicate that a stress change occurred at the depth of the Pacific plate, because the 2011 Tohoku EQ moved the Pacific plate downward at the shallower part, and compressed the deeper part of the plate.

Acknowledgements

We thank captains and crew of KH09-3 cruise, Shincho-maru cruise and aftershock observation cruise of the 2011 Tohoku Earthquake. We used waveform data from Hi-net by NIED.