
[JJ] Evening Poster | S (Solid Earth Sciences) | S-SS Seismology

[S-SS15]Fault Rheology and Earthquake Physics

convener:Hideki Mukoyoshi(Department of Geoscience Interdisciplinary Graduate School of Science and Engineering, Shimane University), Wataru Tanikawa(Japan Agency for Marine-Earth Science and Technology, Kochi Institute for Core Sample Research), Takanori Matsuzawa(国立研究開発法人 防災科学技術研究所, 共同), Keisuke Yoshida(Tohoku University)

Mon. May 21, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

The goal of this session is to integrate theoretical, experimental, observational, and numerical perspectives from various fields such as seismology, geodesy, geology, mineralogy, and so on, to define what is known about earthquake source processes and the physical and chemical elementary processes of faulting. This session welcomes studies that address such issues as pre-, co-, and post-seismic processes, the rheology of seismogenic faults and fault rocks, laboratory experiments on elementary processes, numerical models based on frictional laws, and estimates of the stress field in the seismogenic zones. We also welcome studies on fault-zone drilling projects and in situ stress measurements.

[SSS15-P09]Effect of periodic loading on the repeating behavior of short-term slow slip events

*Takanori Matsuzawa¹, Yoshiyuki Tanaka², Bunichiro Shibazaki³ (1.National Research Institute for Earth Science and Disaster Resilience, 2.Earthquake Research Institute, University of Tokyo, 3.Building Research Institute)

Keywords:Slow Earthquakes, Slow Slip Event, Numerical Simulation

It has been reported that Earth tides affect the activity of episodic tremor and short-term slow slip events (hereinafter, short-term SSEs) in Nankai and Cascadia (e.g., Nakata et al., 2008; Rubinstein et al., 2008; Yabe et al., 2015). In addition, Ide and Tanaka (2014) reports that long-period loading by the ocean also affects the occurrence of deep low frequency tremor. The tidal effect on the SSEs has been also examined by numerical studies (e.g., Hawthorne and Rubin, 2013). In our previous study, we considered the stress perturbation on the plate interface by the Earth tides (especially, semi-diurnal (M2) and fortnight (Mf) tides). In this study, we examined the case with the longer period which is dominant in ocean loading.

We evaluated the effect of periodic loading assuming a flat plate interface with the dipping angle of 15 degrees. Our numerical model is similar to our previous study (Matsuzawa et al., 2010). A rate- and state-dependent friction law (RS-law) with cutoff velocities is adopted as the friction law on each element. We assume a circular short-term SSE region with the radius of 6 km at the depth of 32 km. (a-b) value in the RS-law is negative within the short-term SSE region, and positive outside the region. Stress perturbation is given by sine function with the amplitude of 0.1 kPa. Incorporating this stress perturbation, we calculate the evolution of slip on the plate interface.

In the case of the period of 1 year, which is dominant in actual ocean loading, the recurrence interval of SSEs becomes shorter than that of the previous one around the maximum of stressing rate. On the other hand, the recurrence interval becomes longer around the minimum of stressing rate. These are similar to the case with shorter loading period (e.g., M2 tides). However, the recurrence of SSEs is not clear for about eight years after the large slip at surrounding area. This is similar to the case without periodic loading, while recovery of repeating SSEs can be seen after one year in the case with M2 and Mf tides.

Such recovery gradually becomes quicker with the decrease of the period from 1year.

Acknowledgment: This work was partially supported by JSPS KAKENHI Grant Numbers JP26800240.