

[JJ] Evening Poster | S (Solid Earth Sciences) | S-VC Volcanology

## [S-VC41]Active Volcanism

convener:Yuta Maeda(Nagoya University), Takahiro Miwa(National research institute for earth science and disaster prevention), Yosuke Aoki(東京大学地震研究所, 共同), Takeshi Nishimura(Department of Geophysics, Graduate School of Science, Tohoku University)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

This session discusses various aspects of active volcanisms including, but not limited to, recent and historical eruptions, various phenomena associated with the volcanic activities, underground structures of the volcanoes, and developments of new instruments based on geophysical, geochemical, geological, and multidiscipline approaches. We also welcome studies on understanding and predicting the transitions of the eruptive activities from observational, theoretical, and experimental approaches.

## [SVC41-P32]Establishment of tephra database in and around the Izu

### Islands &ndash; a case of the Daisan-yama site on Niijima

\*Kaori Aoki<sup>2</sup>, Makoto Kobayashi<sup>2</sup>, Fumikatsu NISHIZAWA<sup>2</sup>, Takehiko Suzuki<sup>1,2</sup> (1.Faculty of Urban Environmental Sciences, Tokyo Metropolitan University, 2.Research Center for Volcanic Hazards and Their Mitigation, Tokyo Metropolitan University)

Tokyo Metropolis has the twenty-three wards, the Tama district, and the islands area. The population of the islands area is 25,695 (1st January 2018). The islands area, namely, the Izu-Ogasawara Islands consist of many volcanic islands and seamounts, excluding some islands. For residents in this area, volcanic disasters are always the threat. Accordingly, to integrate the fundamental information for preventing volcanic disasters, we investigate the history of volcanic activities, the pattern of each eruption, and the mass volume of erupted tephtras and try to establish the high-resolution tephrochronology (Kobayashi *et al.*, 2018; Nishizawa *et al.*, 2018; Ito *et al.*, 2018, this session).

The Izu-Ogasawara islands is the islands arc which is approximately 1100 km long and 300 km to 400 km wide. The Sagami trough off the southeastern Izu Peninsula is connected to the Ogasawara Trench to the southward. This island arc consists of the Ogasawara ridge running parallel to the west of the Ogasawara trench, westly the Sichito-Iwoujima ridge across the Ogasawara trough, and more westly the Nishi-Shichito ridge on the east of the Shikoku basin. The most active volcanic belt belongs on the Sichito-Iwoujima ridge, and the northernmost Izu-Oshima is located at 110 km south from the Tokyo Metropolitan Government building, the southernmost Minami-Iwoujima is located at 1281 km south. The north part of the Nishi-Shichito ridge has the several seamount chains, and the northernmost and largest chain is the Zenisu ridge. Next, the summary on the location of seven main volcanic islands belonging to the Izu Islands is as follows. Izu-Oshima, Niijima, and Kozushima are located on the Zenisu ridge from east to southwest. Furthermore, from Izu-Oshima, Toshima is located 25 km southwest, Miyakejima is 72 km south, Mikurajima is 96 km south, and Hachijyojima is 179 km south. Four islands, Izu-Oshima, Toshima, Niijima and Kozushima, are located within 60-km radius, and six islands adding Miyakejima and Mikurajima somewhat southward to above four islands are located within 100-km radius. Tephtras provided by the large eruptions from each island and volcanos on the Japanese mainland would spread across the sea and deposit on the plural islands. Such tephtras reported in Sugihara *et al.* (2005) and Saito *et al.* (2006) have been useful as the key bed to correlate the stratigraphy between islands. On studying tephrostratigraphy in and around islands, it is difficult to try tracking survey on the distribution and thickness of each tephtra in the distance, therefore the tephtra correlation is on the basis of the petrographic description and geochemical data. This project focuses on establishing the database of tephtras collected form each island of the Izu Islands and their petrographic

description and geochemical data. Also, it was likely that such tephra across the sea would deposit in the hemipelagic sediment around volcanic islands. Our tephra database of the Izu Islands should be a powerful tool for marine tephra correlations in future. Today, we present the result of tephra analysis at the Daisan-yama site on Niijima as the part of on-going database project.

At the southwestern slope of the Daisan-yama in Niijima, we can find at least seventeen visible tephra layers, consisting of the tephra including Fujimi-pass tephra group from Niijima, the tephra from Kozushima, and thin basaltic tephra (Kobayashi *et al.*, 2018). After tephra samples were washed by the ultrasonic cleaner, we collected the description of petrographic characters, major-element chemistry determined by EDS and refractive indices of volcanic glass shards in each tephra. Also, AT and K-Ah tephra were detected as cryptotephra from loam samples. Tephra records during the last 30 ka at this site is one of the important information to construct the high-resolution chronology in the area.