

[JJ] Evening Poster | S (Solid Earth Sciences) | S-VC Volcanology

## [S-VC41]Active Volcanism

convener:Yuta Maeda(Nagoya University), Takahiro Miwa(National research institute for earth science and disaster prevention), Yosuke Aoki(東京大学地震研究所, 共同), Takeshi Nishimura(Department of Geophysics, Graduate School of Science, Tohoku University)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

This session discusses various aspects of active volcanisms including, but not limited to, recent and historical eruptions, various phenomena associated with the volcanic activities, underground structures of the volcanoes, and developments of new instruments based on geophysical, geochemical, geological, and multidiscipline approaches. We also welcome studies on understanding and predicting the transitions of the eruptive activities from observational, theoretical, and experimental approaches.

## [SVC41-P47]Average magma ascent rate and xenolith-entrained depth constrained by thermal effect on crustal xenoliths

\*Hiroshi Kawabata<sup>1</sup>, Koshi Nishimura<sup>2</sup>, Kenji Horie<sup>3</sup>, Mami Takehara<sup>3</sup> (1.Faculty of Science and Technology, Kochi Univeristy, 2.Natural Science Laboratory. Toyo University, 3.National Institute of Polar Research)

Keywords:xenolith, partial melting, thermal history

Xenoliths are one of the important sources for clues on crustal structure and magma ascent rate. However, determination of xenolith-entrained depth is not straightforward especially for thermally affected xenoliths, in which mineral assemblage and compositions have been modified. In addition, commonly used Stokes's settling law only indicates the minimum magma ascent rate required to transport xenoliths to the surface. To overcome some of the limitations, we considered the timescale of heat transfer and disequilibrium phenomena of xenolith melting. We performed a non-steady-state 1-D thermal conduction model in spherical coordinates, varying initial temperature conditions and xenolith size. The model involves latent heat of fusion and temperature-dependence of thermal conductivity. Boundary conditions specified temperature, where xenolith surface has constant temperature as the host magma. This implicitly assumes that the xenoliths are volumetrically insignificant. Melt productivity in xenolith, which is also essential in the calculation, is based on phase equilibria but the effect of disequilibrium melting is also considered. Therefore, melt proportions at a given position increase by increasing both the degree of superheating and the heating time until melt proportions reach the values expected from phase equilibria. As an example, the above model was applied for Cretaceous granitic xenoliths entrained in a Miocene intermediate dike in NE Shikoku, SW Japan. Unknown initial xenolith temperature ( $T_0$ ) was estimated using a xenolith entrained-depth and a geotherm gradient for shallow crust (30°C/km). For xenoliths with 10–60 cm in diameter and  $T_0$  of 40–460°C enclosed by a host magma with 900°C, it took ~0.3–12 hours to reach solidus temperature at the xenolith center. If disequilibrium melting is assumed, longer time (order of days) is required for 5% melting at the xenolith center. Based on the resultant timescale, we can constrain the combination of average magma ascent rate and entrained-depth that explains the observed xenolith texture (i.e., the presence or absence of glass at the center of a xenolith). The approach proposed here would be applicable for other crustal xenoliths as long as the xenoliths were entrained in ascending magmas.