

[JJ] Evening Poster | S (Solid Earth Sciences) | S-VC Volcanology

[S-VC44]Magmatism and volcanic dynamics on subduction zone

convener:Yujiro Suzuki(Earthquake Research Institute, The University of Tokyo), Hitomi Nakamura(Department of Solid Earth Geochemistry, Japan Agency for Marine-Earth Science and Technology), Yu Iriyama(国立研究開発法人 防災科学技術研究所)

Thu. May 24, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Subduction zones are characterized by magmatism and volcanism caused by magma generation, magma transportation and storage, and volcanic eruption. These magmatic process and volcanic dynamics have been investigated using laboratory experiments and sophisticated numerical modelings. Recent progress of observation techniques has revealed the structures under volcanoes. Besides, the studies based on seismological observations and geochemical analyses try to determine the spatial and temporal scale of magmatic activity and explain the process of magma generation. This session aims to discuss the latest studies about magmatism and volcanic dynamics from various perspectives and integrate these understandings.

[SVC44-P04]Petrological study of the Miocene adakitic andesite in the Tateyama area, eastern part of Toyama prefecture.

*Watanabe Syun¹, Toshiro Takahashi¹ (1.Niigata university)

Keywords:Miocene , Hokuriku district, adakite

The Oligocene ~ Miocene volcanic rocks are widely distributed in Hokuriku Green Tuff Province. Some previous studies discussed that magma generation and evolution process in the Hokuriku district at Oligocene ~ Miocene (e.g. Uematsu et al., 1995; Takahashi and Shuto, 1999; Ishiwatari and Ohhama, 1997). However, tectonic setting in Hokuriku district at Miocene remains obscure in detail.

The Iwaine formation (Early ~ Middle Miocene) and the Fukuhira formation (Middle Miocene) consist of mainly andesitic pyroclastic rocks and lavas.

The Miocene volcanic rocks in this study area can be divided into four groups by petrographic and chemical features (Group-1; tholeiitic low-K basalt, Group-2: tholeiitic andesite, Group-3; calc alkaline amphibole andesite, Group-4; calc-alkaline two-pyroxene andesite). The Iwaine formation consist of calc-alkaline rocks (group-3, group-4) and tholeiitic rocks (group-1, group2). Fukuhira formation consist of tholeiitic rocks (group-3, group-4).

Especially, calc-alkaline amphibole andesite (Group-3) is characterized by relatively higher Sr/Y ratio contents than common andesite in island-arc. These chemical features of Group-3 is similar to adakite that is generated by partial melting of subducted oceanic plate (e.g. Defant and Durmond, 1990). This study aimed to discussion of the magma generation and evolution process of Group-3 andesite and unravels the Miocene magmatism in the Hokuriku district.

Phenocryst assemblage of group-3 is amphibole + plagioclase + clinopyroxene + orthopyroxene. Almost phenocrystic plagioclases have dusty zone and honeycomb texture. Phenocrystic clinopyroxene and orthopyroxene in the Group-3 andesite can be divided into three types by petrographic features (Type-1; monocrystal pyroxene, Type-2; glomeroporphyritic texture with amphibole, Type-3; glomeroporphyritic texture with pyroxene). Type-1 pyroxene has narrow range Mg# (0.77~0.80), Type-2 pyroxene has wide range Mg# (0.70~0.86), Type-3 pyroxene has wide range Mg# (0.68~0.80).

Group-3 is plotted in calc-alkaline field on the FeO*/MgO -SiO₂ diagram (Miyashiro, 1974), and low-K andesite field on the SiO₂-K₂O diagram (Gill, 1981). In term of Y-Sr/Y diagram (Defant and Durmond, 1990), group-3 plotted in overlap field between adakite and island arc-ADRs. However, Group-3 is

characterized by higher MgO, Cr and Ni content than the typical adakite.

Petrographical and chemical compositions of phenocrystic minerals in Group-3 andesite exhibit disequilibrium texture and compositions such as dusty zoned plagioclase and coexistence of high Mg# pyroxene and low Mg# pyroxene. Thus, these petrographical and petrological features suggest that magma evolution process of Group-3 andesite involves magma mixing.

High Mg# (Mg# 0.85~0.86) pyroxene shows Type-2 texture, and is equilibrium with high-magnesian andesite melt such as Oligocene high-magnesian andesite in Noto peninsula (Uematsu et al., 1995). Middle Mg# (Mg# 0.77~0.83) pyroxene shows Typ-1, Type-2, Type3 texture, and is equilibrium with whole rock composition of group-3. Low Mg# (Mg# 0.68~0.74) pyroxene show Type1, Type-2 texture, and is equilibrium with dacitic ~ rhyolitic melt such as primitive slab melt (Zamora, 2000).

Mg# of phenocrystic pyroxene indicate that Group-3 magma is generated by magma mixing between mafic end member magma (HMA or bronzite-andesite) and felsic end member magma. Whole rock composition of Group-3 andesite indicates possibility of felsic end member is adakite magma.

Therefore, it is thought that Group-3 andesite generated by magma mixing between high-magnesian andesite magma and dacitic ~ rhyolitic magma. Additionally, the chemical composition of felsic end-member magma needs high Sr/Y ratio such as adakite magma. It is thought that these results give limitation to discuss about the tectonic setting model of Miocene magmatism in the Hokuriku green tuff province.