

河畔林の発達した河川（櫛田川派川・祓川，三重県）におけるシジミ2種
（ヤマトシジミ，マシジミ）の炭素・窒素同位体比と餌資源

Carbon and nitrogen isotope ratios and diets of *Corbicula japonica*
and *Corbicula leana* in the Harai River (a branch river of the Kushida
River, Mie Prefecture, central Japan) with rich riparian forests

牛川 晃¹、*杉谷 健一郎¹、山本 真里子¹、山本 鋼志¹、村岡 一幸⁴、北村 淳一³、太田 民久⁵、原口 岳²、
陀安 一郎²

Akira Ushikawa¹, *Kenichirio Sugitani¹, Mariko Yamamoto¹, Koshi Yamamoto¹, Kazuyuki Muraoka⁴,
Jyunichi Kitamura³, Tamihisa Ohta⁵, Takashi Haraguchi², Ichiro Tayasu²

1. 名古屋大学、2. 総合地球環境学研究所、3. 三重県総合博物館、4. 三重県立松阪高校、5. 富山大学

1. Nagoya University, 2. Research Institute for Humanity and Nature, 3. Mie Prefectural Museum, 4. Matsusaka High School, 5. University of Toyama

In order to eventually reveal factors controlling distribution and abundance of the bivalve *Unionidae* group in the Harai River, a branch of the Kushida River, Mie Prefecture, central Japan, the bivalve *Corbicula* as alternatives were analyzed for carbon and nitrogen isotope ratios. The Harai River is withdrawn from a dam constructed at the Kushida River, and has rich riparian forests. Samples collection was performed 4 times in 2016 and 2017, at 4 localities; *C. leana* from the two upstream and midstream sites (Sites 1 and 11) and *C. japonica* from the two estuary sites (Sites 15 and 16). Particulate organic matter (POM) in river water was also collected from 5 sites including those for *Corbicula* collections. We obtained 54 and 58 isotopic data for *C. leana* and for *C. japonica*, respectively, and 20 for POM. While carbon and nitrogen isotope ratios of *Corbicula* range widely from -26.3 to -22.2 per mil and from 5.6 to 10.8 per mil, respectively, samples from each site cluster closely (Figure). Seasonal variation is negligible, except for the *C. leana* of Site 11, compared differences between sites. Two populations of *C. leana* and one population of *C. japonica* comprise an array of negative correlation between carbon and nitrogen isotope ratios, which is parallel to the distribution of POM. Carbon and isotopic ratios of these three *Corbicula* populations are likely constrained by averaged available food sources (POM) at each site. Population of *C. japonica* from the lowermost site shows a positive correlation between carbon and nitrogen isotope ratios, implying their utilization of two end-members of POMs with distinct carbon and nitrogen isotopic ratios. *C. japonica* samples in this study was significantly lower in carbon and nitrogen isotope ratios than those reported by Kasai and Nakata (2005) (Fish. Sci. 71, 151-158), who demonstrated that terrestrial organic matter was significantly important even for *C. japonica* diet. The distinct result of this study may be explained by richness of riparian forests at Harai River, which have made *C. japonica* there further depend on terrestrial food sources. On the other hand, *C. leana* population from the uppermost site of the Harai River has heaviest carbon isotopic ratios, which could be attributed to an increase in utilization of planktons flourished in the dam.

Reference

Kasai, A. and Nakata, A. (2005) Utilization of terrestrial organic matter by the bivalve *Corbicula japonica* estimated from stable isotope analysis. Fisheries Science 71, 151-158.

キーワード：シジミ、同位体比、餌資源、陸起源有機物、祓川

Keywords: Corbicula, isotopic ratios, food sources, terrestrial organic matter, Harai River

