

Characteristics of radiocesium-bearing microparticles and Type B particles using multiple synchrotron radiation X-ray analyses

*三浦 輝^{1,2}、栗原 雄一²、山本 政義³、山口 紀子⁴、坂口 綾⁵、桧垣 正吾⁶、高橋 嘉夫²

*Hikaru Miura^{1,2}, Yuichi Kurihara², Masayoshi Yamamoto³, Noriko Yamaguchi⁴, Aya Sakaguchi⁵, Shogo Higaki⁶, Yoshio Takahashi²

1. 電力中央研究所 環境科学研究所、2. 東京大学大学院 理学系研究科、3. 金沢大学 LLRL、4. 農研機構、5. 筑波大学 アイソトープ環境動態研究センター、6. 東京大学 アイソトープ総合センター

1. Environmental Science Research Laboratory, Central Research Institute of Electric Power Industry, 2. Department of science, Graduate school of The university of Tokyo, 3. Low Level Radioactivity Laboratory, University of Kanazawa, 4. Institute for Agro-Environmental Sciences, NARO, 5. Center for Research in Isotopes and Environmental Dynamics, University of Tsukuba, 6. Isotope Science Center, The University of Tokyo

Introduction: A large amount of radiocesium was emitted into environment by the Fukushima Nuclear Power Plant (FDNPP) accident in March, 2011. Adachi et al. (2013) reported glassy water-insoluble microparticles including radiocesium, called as radiocesium-bearing microparticles (CsMPs). The CsMP is spherical with 1-3 μm in diameter and the radioactivity ranges from 0.5 to 4 Bq. It has been suggested that the CsMP was mainly emitted from Unit 2 or Unit 3 of FDNPP based on the $^{134}\text{Cs}/^{137}\text{Cs}$ activity ratio in the samples. In contrast, Ono et al. (2017) reported new particles called as Type B particles emitted from Unit 1. Type B particles are in various shapes and the size is 50-300 μm with radioactivity ranging from 30 to 100 Bq. These differences may represent the difference of generating process or condition of each unit in the plant. Previous studies have reported chemical properties of radioactive particles in detail but the number of particles reported is small. In this study, we tried to understand radioactive particles systematically by analyzing a lot of particles separated using wet separation method.

Method: In this study, we collected 53 Type B particles and 13 CsMPs from road dusts, non-woven fabric cloths from Fukushima, and aerosol filters from Kanagawa Prefecture by a wet separation method. After measurement of radioactivity with a high-purity germanium semiconductor detector, scanning electron microscope and energy dispersive X-ray spectroscopy analyses were performed to confirm that separated particles were CsMPs or Type B particles. We investigated inner structure and calculated the volume and porosity of Type B particles by X-ray μ -computed tomography (CT). We determined Rb/Sr ratio by X-ray fluorescence (XRF) analysis. Redox condition of each unit was investigated by X-ray adsorption near edge structure (XANES) analysis for Uranium in particles.

Result: CT combined with XRF analysis showed the presence of many voids and iron particles in Type B particles. In addition, ^{137}Cs concentration of CsMPs were ~ 10000 times higher than that of Type B particles, which suggests that Type B particles were formed by fuel melt. In contrast, CsMPs were formed by gas. Among Type B particles, spherical particles had higher ^{137}Cs concentration than non-spherical particles. Type B particles with larger porosity had higher ^{137}Cs radioactivity because of capturing a lot of volatile elements such as Cs and Rb within the particles. Moreover, four spherical particles had inclusions in their voids which are considered to be formed by rapid cooling of gaseous materials. XANES analysis showed the presence of U(IV) in a Type B particle, whereas U(VI) in other Type B particles and a CsMP.

These results suggest that Type B particles and CsMPs are totally different in forming process and they have information of condition in Units.

キーワード：放射性粒子、セシウム、X線分析

Keywords: Radioactive particle, Cesium, X-ray analyses