## Factors controlling the distribution of ferromanganese nodules in the Minamitorishima EEZ

\*中村 謙太郎<sup>1</sup>、下村 遼<sup>1</sup>、堀之内 航一<sup>1</sup>、町田 嗣樹<sup>2,3</sup>、安川 和孝<sup>1,2</sup>、藤永 公一郎<sup>2,1</sup>、加藤 泰浩<sup>1,2,3</sup> \*Kentaro Nakamura<sup>1</sup>, Ryo Shimomura<sup>1</sup>, Koichi Horinouchi<sup>1</sup>, Shiki Machida<sup>2,3</sup>, Kazutaka Yasukawa <sup>1,2</sup>, Koichiro Fujinaga<sup>2,1</sup>, Yasuhiro Kato<sup>1,2,3</sup>

 1. 東京大学大学院工学系研究科システム創成学専攻、2. 千葉工業大学次世代海洋資源研究センター、3. 海洋研究開発機構
1. Department of Systems Innovation, School of Engineering, University of Tokyo, 2. ORCeNG, Chiba Institute of Technology, 3. JAMSTEC

Ferromanganese nodules are polymetallic concretions consisting mainly of manganese and iron, which are found in deep-sea surficial sediments in many areas of the world ocean [1]. Since 1960s, the nodules are of high economic interest due to their enrichment of valuable metals such as Ni, Cu, and Co [2]. Very recently, large-scale distribution of ferromanganese nodules has also been confirmed within Japanese exclusive economic zone (EEZ) around Minamitorishima island [3, 4] and, for future exploitation of the nodules in the Minamitorishima EEZ, detailed investigation to elucidate the origin and distribution of the nodules is needed.

Ferromanganese nodules have long been studied extensively, especially those occurring between the Clarion and Clipperton fracture zones [5]. However, many important questions still remain unanswered regarding the formation and distribution of the nodules. In particular, factors controlling the distribution of ferromanganese nodules have been one of the most important problems to solve. Correlation between nodule distribution and deep-sea currents has long been pointed out in the southern Pacific and southern Indian Oceans [6, 7]. Moreover, it has been considered that hiatus of the underlying sediments, caused by deep-sea currents, is also related to the occurrence of ferromanganese nodules [6, 8].

In this presentation, we will discuss the factors controlling the distribution of ferromanganese nodules in the Minamitorishima EEZ in relation to underlying sediments and deep-sea currents.

## References:

Petersen et al. (2016) *Marine Policy*, **70**, 175–187, [2] Hein et al. (2013) *Ore Geology Reviews*, **51**, 1–14, [3] Machida et al. (2016) *Geochemical Journal*, **50**, 539–555, [4] JAMSTEC press release, August 26, 2016, [5] ISA (2012) Briefing Paper 01/12, [6] Kennett and Watkins (1975) *Science*, **188**, 1011–1013, [7] Wright et al. (2005) *New Zealand Journal of Geology and Geophysics*, **48**, 27–41, [8] Usui et al. (1987) *Marine Geology*, **74**, 237–275.

## キーワード:マンガンノジュール、南鳥島EEZ、分布、直下堆積物、深層海流

Keywords: ferromanganese nodules, Minamitorishima EEZ, distribution, underlying sediments, deep-sea currents