

A Spatially Resolved AU-scale Inner Disk around DM Tau with ALMA

*橋本 淳¹、Liu Hauyu⁷、Dong Ruobing⁶、小西 美穂子²、武藤 恭之³、長谷川 靖紘⁴、塚越 崇⁵、工藤 智幸²

*Jun Hashimoto¹, Hauyu Liu⁷, Ruobing Dong⁶, Mihoko Konishi², Takayuki Muto³, Yasuhiro Hasegawa⁴, Takashi Tsukagoshi⁵, Tomoyuki Kudo²

1. 自然科学研究機構アストロバイオロジーセンター、2. 自然科学研究機構国立天文台、3. 工学院大学、4. ナサ ジェット推進研究所、5. 茨城大学、6. アリゾナ大学、7. マックスプランク天文学研究所

1. The National Institutes of Natural Sciences, Astrobiology center, 2. The National Institutes of Natural Sciences, National Astronomical Observatory of Japan, 3. Kogakuin University, 4. NASA Jet Propulsion Laboratory, 5. Ibaraki University, 6. Arizona University, 7. Max Planck Institute for Astronomy

We present Atacama Large Millimeter/submillimeter Array (ALMA) observations of the dust continuum at 1.3 mm and 12CO (2-1) line of the transitional disk around DM Tau. DM Tau exhibits no near-infrared (NIR) excess in its spectral energy distribution (SED), which means the inner cavity is well depleted. However we found a spatially resolved inner disk at about $r = 3$ AU in the dust continuum image. Assuming the inner disk's temperature of 200 K, a dust mass of the inner disk is about 0.2 MJup. The brightness structures in both the dust continuum and the 12CO (2-1) are marginally asymmetric: 1.15 +/- 0.08 and 1.47 +/- 0.16 times brighter in the northwest part in the inner disk, respectively. We also performed a simple analytic modeling of the disk's brightness profile, and derived physical disk parameters by fitting the observed visibilities. Our modeling efforts found DM Tau's dust disk consists of three components: an inner disk, an outer disk, and an extended structure. In the presentation, we will discuss possible origins of the multi-ring structure around DM Tau.

キーワード：原始惑星系円盤、アルマ、遷移円盤

Keywords: protoplanetary disk, ALMA, transitional disk