

Numerical Simulation of Traveling Ionospheric Disturbances generated by upward propagating gravity waves

*三好 勉信¹、陣 英克²、藤原 均³、品川 裕之²

*Yasunobu Miyoshi¹, Hidekatsu Jin², Hitoshi Fujiwara³, Hiroyuki Shinagawa²

1. 九州大学大学院理学研究院地球惑星科学部門、2. 情報通信研究機構、3. 成蹊大学

1. Kyushu University, 2. NICT, 3. Seikei University

It has been recognized that gravity waves (GWs) play an important role on the momentum and energy budget in the thermosphere/ionosphere. In this study, using a whole atmosphere-ionosphere coupled model (GAIA) with a high horizontal resolution, behaviors of Traveling Ionospheric Disturbances (TIDs) generated by upward propagating GWs in the thermosphere are investigated. The horizontal resolution of GAIA is 1 degree longitude by 1 degree latitude, which is adequate to simulate large-scale GWs. The GAIA contains the region from the ground surface to the upper thermosphere, so that we can simulate excitation of gravity waves in the lower atmosphere, their upward propagation to the mesosphere and thermosphere, and their impacts on the thermosphere/ionosphere system. The GAIA can simulate TIDs because interaction processes between the ionosphere and neutral atmosphere are included. The equatorward (poleward) TIDs become dominant during the daytime (nighttime) due to the filtering effect of GWs by poleward (equatorward) thermospheric winds. The dominant horizontal wavelength and period of the simulated TIDs are 500–1500 km and 30–90 min, respectively. The dominant wavelength and period of TIDs are the same as those of GWs near 250–300 km height. We discuss seasonal and longitudinal variations of TIDs and their relation to GW activity in the thermosphere.

キーワード：電離圏—大気圏結合過程、伝播性電離圏擾乱、大気重力波

Keywords: ionosphere-atmosphere coupling process, TID, Gravity wave