

Ionospheric responses to the 21 August 2017 Solar Eclipse by using data assimilation approach

*陳 佳宏¹、Lin Charles¹、Matsuo Tomoko²

*Chia-Hung Chen¹, Charles Lin¹, Tomoko Matsuo²

1. 台湾国立成功大学地球科学学科、2. コロラド大学航空宇宙工学

1. Department of Earth Sciences, National Cheng Kung University, 2. Ann and H.J. Smead Department of Aerospace Engineering Sciences, University of Colorado at Boulder, CO

Using the physical based thermosphere-ionosphere model with ensemble Kalman filter, this study reports the first data assimilation results of the ionosphere responses to the solar eclipse on 21 August 2017. The system with 2-minute assimilation cycle of ground-based GNSS observations show the dynamic variations of the equatorial ionization anomaly (EIA) due to modifications of the electrodynamics by the solar eclipse. Two major ionosphere responses are captured. First, there was an early appearance of EIA at the westward boundary of moon shadow, followed by the feature of the enhanced EIA at lower latitudes and the suppressed EIA at the higher latitudes. These eclipse-induced conjugate EIA variations are produced by the eastward electric field perturbation around magnetic equator and the westward electric field perturbation at the higher latitudes.

キーワード：電離圏データ同化モデル、データ同化による電場変化、日食による電離圏変化

Keywords: ionospheric data assimilation model, electric field, solar eclipse-induced ionospheric responses