

## Electron Scattering by Chorus Waves Generating Pulsating Aurora

\*Satoshi Kasahara<sup>1</sup>, Yoshizumi Miyoshi<sup>2</sup>, Shoichiro Yokota<sup>3</sup>, Takefumi Mitani<sup>4</sup>, Yoshiya Kasahara<sup>5</sup>, Shoya Matsuda<sup>2</sup>, Atsushi Kumamoto<sup>6</sup>, Ayako Matsuoka<sup>4</sup>, Yoichi Kazama<sup>7</sup>, Harald U Frey<sup>8</sup>, Vassilis Angelopoulos<sup>9</sup>, Satoshi Kurita<sup>2</sup>, Kunihiro Keika<sup>1</sup>, Kanako Seki<sup>1</sup>, Iku Shinohara<sup>4</sup>

1. The University of Tokyo, 2. Nagoya University, 3. Osaka University, 4. ISAS, 5. Kanazawa University, 6. Tohoku University, 7. ASIAA, 8. UCB, 9. UCLA

Pulsating aurorae, which are quasiperiodic, blinking patches of light tens to hundreds of kilometres across, appear at altitudes of about 100 kilometres in the high-latitude regions of both hemispheres, and multiple patches often cover the entire sky. This auroral pulsation, with periods of several to tens of seconds, is generated by the intermittent precipitation of energetic electrons (several to tens of kiloelectronvolts) arriving from the magnetosphere and colliding with the atoms and molecules of the upper atmosphere. A possible cause of this precipitation is the interaction between magnetospheric electrons and electromagnetic waves called whistler-mode chorus waves. However, no direct observational evidence of this interaction has been obtained so far. Here we report that energetic electrons are scattered by chorus waves, resulting in their precipitation. Our observations were made in March 2017 with a magnetospheric spacecraft equipped with a high-angular-resolution electron sensor and electromagnetic field instruments. The measured quasiperiodic precipitating electron flux was sufficiently intense to generate a pulsating aurora, which was indeed simultaneously observed by a ground auroral imager.

Keywords: pitch angle scattering, chorus waves, ERG (Arase)