

Initial results of EMIC observation by MGF/Arase

*野村 麗子¹、松岡 彩子²、寺本 万里子³、三好 由純³、松田 昇也³、栗田 怜³、桂華 邦裕⁴、山本 和弘⁵、能勢 正仁⁵、小路 真史³、藤本 晶子⁶、篠原 学⁷、田中 良昌⁸、笠原 禎也⁹、笠羽 康正¹⁰、石坂 圭吾¹¹

*Reiko Nomura¹, Ayako Matsuoka², Mariko Teramoto³, Yoshizumi Miyoshi³, Shoya Matsuda³, Satoshi Kurita³, Kunihiro Keika⁴, Kazuhiro Yamamoto⁵, Masahito Nose⁵, Masafumi Shoji³, Akiko Fujimoto⁶, Manabu Shinohara⁷, Yoshimasa Tanaka⁸, Yoshiya Kasahara⁹, Yasumasa Kasaba¹⁰, Keigo Ishisaka¹¹

1. 宇宙航空研究開発機構、2. 宇宙科学研究所、3. 名古屋大学宇宙地球環境研究所、4. 東京大学、5. 京都大学、6. 九州大学、7. 鹿児島工業高等専門学校、8. 国立極地研究所、9. 金沢大学、10. 東北大学、11. 富山県立大学

1. Japan Aerospace and Exploration Agency, 2. ISAS, 3. ISEE, 4. The University of Tokyo, 5. Kyoto University, 6. Kyusyu University, 7. Kagoshima National College of Technology, 8. National Institute of Polar Research, 9. Kanazawa University, 10. Tohoku University, 11. Toyama Prefectural University

The Electromagnetic ion cyclotron (EMIC) wave is one of the key phenomena to understand the loss dynamics of MeV-energy electrons in the Earth's radiation belt. The precipitation into the ionosphere of MeV-energy electrons was observed in the isolated proton aurora which is an ionospheric footprint for the region of EMIC wave-particle interaction in the magnetosphere [Miyoshi+2008]. However, it is still unclear how significant the EMIC wave is in terms of the global scale of the whole radiation belt, and it is important to understand where and when the preferable condition is set in the radiation belt of the inner magnetosphere for strong EMIC waves.

Li et al. [2011a, 2011b] showed that the whistler mode chorus waves, as a counter part of EMIC waves, were modulated by the compressional ULF waves and the variation of background density, and showed the global distribution of its modulation with three different mechanism, by compressional ULF waves, by density depletions, and by density enhancement. In order to find the global distribution of EMIC wave modulation and its mechanism, we investigated in situ statistical survey of EMIC waves observed by Arase satellite from March to October 2017, and identified 64 events using the fluxgate magnetometer (MGF) on the satellite. We found that the enhancements of EMIC waves were occurred at localized region less than 0.1MLT, and with 5-10nT weaker magnetic field than background, at 22:20-22:30UT on 27 May and at 2017, and at 07:30-08:00UT on 25 June 2017. We will discuss the EMIC wave excitation and enhancement mechanism in the deep inner magnetosphere.