Spatial distribution of radiation belt protons deduced from unexpected output of HEP on board the Arase satellite

*三宅 亙¹、戸田 穂乃香¹、三谷 烈史²、高島 健²、三好 由純³ *Wataru Miyake¹, Honoka Toda¹, Takefumi Mitani², Takeshi Takashima², Yoshizumi Miyoshi³

1. 東海大学工学部、2. JAXA宇宙科学研究所、3. 名古屋大学宇宙地球環境研究所

1. Tokai University, 2. ISAS/JAXA, 3. ISEE/Nagoya University

HEP instrument on board the Arase satellite measures electrons with energies from 70 keV to 2 MeV. The Van Allen probes observation revealed that the electrons of MeV energy range never exist in the inner radiation belt. However, the HEP has detected unexpected counts even at the 2 MeV energy channel in the inner radiation belt (L < 2.0). Any contamination of high energy protons (> 30 MeV) is supposed to result in the unexpected counts. By assuming no contribution from electrons, we can obtain spatial distribution of energetic protons from the unexpected counts. The result shows a concentration on the magnetic equator especially at L > 1.5, which is basically consistent with pitch angle observation of tens of MeV protons near the magnetic equator from the Van Allen probes.

キーワード:プロトン放射線帯、あらせ衛星 Keywords: proton radiation belt, Arase satellite