

## Volcanic Control of Jupiter's Aurora and Middle Magnetosphere Dynamics Observed by Hisaki/EXCEED

\*Chihiro Tao<sup>1</sup>, Tomoki Kimura<sup>2</sup>, Fuminori Tsuchiya<sup>3</sup>, Go Murakami<sup>4</sup>, Kazuo Yoshioka<sup>5</sup>, Atsushi Yamazaki<sup>4</sup>, Sarah V Badman<sup>6</sup>, Hiroaki Misawa<sup>3</sup>, Hajime Kita<sup>3</sup>, Yasumasa Kasaba<sup>3</sup>, Ichiro Yoshikawa<sup>5</sup>, Masaki Fujimoto<sup>4</sup>

1. National Institute of Information and Communications Technology, 2. Riken, 3. Tohoku University, 4. ISAS/JAXA, 5. University of Tokyo, 6. Lancaster University

Temporal variation of Jupiter's northern aurora during enhanced Io volcanic activity was detected using the EXCEED spectrometer on board the Hisaki Earth-orbiting planetary space telescope. It was found that in association with reported Io volcanic events in early 2015, auroral power and estimated field-aligned currents were enhanced during day of year 40–120. Furthermore, the far ultraviolet color ratio decreased during the event, indicating a decrease of auroral electron mean energy and total acceleration by <30%. During the episode of enhanced Io volcanic activity, Jupiter's magnetosphere contains more source current via increased suprathermal plasma density by up to 42%; therefore, it would have required correspondingly less electron acceleration to maintain the enhanced field-aligned current and corotation enforcement current. Sporadic large enhancements in auroral emission detected more frequently during the active period could have been contributed by non-adiabatic magnetospheric energization.

Keywords: Hisaki, Jupiter, Aurora