

## Comparison of visible spectra between Ryugu and low-albedo asteroid families in the inner main belt

\*Seiji Sugita<sup>1</sup>, ERI TATSUMI<sup>1</sup>, Sunao Hasegawa<sup>2</sup>, Suzuki Yudai<sup>1</sup>, Hiroaki Kamiyoshihara<sup>1</sup>, Rie Honda<sup>3</sup>, Shingo Kameda<sup>4</sup>, Tomokatsu Morota<sup>5</sup>, Chikatoshi Honda<sup>9</sup>, Toru Kouyama<sup>8</sup>, Manabu Yamada<sup>7</sup>, Masahiko Hayakawa<sup>2</sup>, yokota yasuhiko<sup>2</sup>, Naoya Sakatani<sup>6</sup>, Hidehiko Suzuki<sup>6</sup>, Kazunori Ogawa<sup>10</sup>, Hirotaka Sawada<sup>2</sup>

1. University of Tokyo, 2. JAXA/ISAS, 3. Kochi University, 4. Rikkyo University, 5. Nagoya University, 6. Meiji University, 7. Chiba Institute of Technology, 8. AIST, 9. University of Aizu, 10. Kobe University

The main mission phase of Hayabusa2 is coming up soon. In this presentation, we will review 1) recent spectroscopic analysis of low-albedo asteroids, particularly in the inner main belt, 2) comparison with Ryugu spectral observations, 3) inflight spectroscopic calibration of our ONC-T.

Keywords: Hayabusa2, asteroids, Ryugu