

Heat conduction in highly porous aggregates

*荒川 創太¹、田中 秀和²、片岡 章雅³、中本 泰史¹

*Sota ARAKAWA¹, Hidekazu Tanaka², Akimasa Kataoka³, Taishi Nakamoto¹

1. 東京工業大学、2. 東北大学、3. 国立天文台

1. Tokyo Institute of Technology, 2. Tohoku University, 3. National Astronomical Observatory of Japan

The thermal conductivity of dust aggregates is a key parameter for many subjects in planetary science (e.g., thermal evolution of small bodies in our solar system). However, the porosity dependence of thermal conductivity for highly porous aggregates is not yet thoroughly understood.

We investigated the thermal conductivity of fluffy dust aggregates. We used the snapshot data of *N*-body simulations of static compression in the periodic boundary condition (Kataoka et al. 2013) and determined the temperature structure and heat flux of the porous dust aggregates using the method of Sirono (2014).

We derived an empirical formula for the thermal conductivity through the solid network as a function of the filling factor of dust aggregates (Arakawa et al. 2017). The results reveal that the thermal conductivity through the solid network is approximately proportional to the square of the filling factor. For the case of dust aggregates with the filling factor in the range of 0.1–0.5, our results are consistent with the experimental data of Krause et al. (2011). For highly porous aggregates whose filling factor is lower than 0.1, it is revealed that the thermal conductivity through the solid network is significantly lower than previously assumed. In light of these findings, we will reexamine the thermal histories of small bodies.

キーワード：彗星、ダストアグリゲイト、熱伝導率、微惑星

Keywords: comet, dust aggregate, thermal conductivity, planetesimal