

Video Observations of Faint Meteors with Tomo-e PM

*大澤 亮¹、酒向 重行¹、猿楽 祐樹²、臼井 文彦³、大坪 貴文⁴、藤原 康德⁵、佐藤 幹也⁶、春日 敏測¹¹、有松 亘⁷、渡部 潤一⁷、土居 守¹、小林 尚人¹、高橋 英則¹、本原 顕太郎¹、宮田 隆志¹、諸隈 智貴¹、小西 真広¹、青木 勉¹、征矢野 隆夫¹、樽澤 賢一¹、森 由貴¹、中田 好一¹、満田 和真¹、一木 真¹、有馬 宣明¹、小島 悠人¹、戸谷 友則⁸、松永 典之⁸、茂山 俊和⁹、板 由房¹⁰、小久保 充¹⁰、前原 裕之¹¹、富永 望¹²、山下 卓也⁷、田中 正臣⁷、猪岡 皓太⁷、池田 思朗¹³、森井 幹雄¹³、吉川 真⁴、浦川 聖太郎¹⁴、奥村 真一郎¹⁴

*Ryou Ohsawa¹, Shigeyuki Sako¹, Yuki Sarugaku², Fumihiko Usui³, Takafumi Ootsubo⁴, Yasunori Fujiwara⁵, Mikiya Sato⁶, Toshihiro Kasuga¹¹, Ko Arimatsu⁷, Jun-ichi Watanabe⁷, Mamoru Doi¹, Naoto Kobayashi¹, Hidenori Takahashi¹, Kentaro Motohara¹, Takashi Miyata¹, Tomoki Morokuma¹, Masahiro Konishi¹, Tsutomu Aoki¹, Takao Soyano¹, Ken'ichi Tarusawa¹, Yuki Mori¹, Yoshikazu Nakada¹, Kazuma Mitsuda¹, Makoto Ichiki¹, Noriaki Arima¹, Yuto Kojima¹, Tomonori Totani⁸, Noriyuki Matsunaga⁸, Toshikazu Shigeyama⁹, Yoshifuksa Ita¹⁰, Mitsuru Kokubo¹⁰, Hiroyuki Maehara¹¹, Nozomu Tominaga¹², Takuya Yamashita⁷, Masaomi Tanaka⁷, Kota Inooka⁷, Shiro Ikeda¹³, Mikio Morii¹³, Makoto Yoshikawa⁴, Seitaro Urakawa¹⁴, Shin-ichiro Okumura¹⁴

1. 東京大学大学院理学系研究科 天文学教育研究センター、2. 京都産業大学 神山天文台、3. 神戸大学 理学研究科、4. 国立研究開発法人宇宙航空研究開発機構 宇宙科学研究所、5. 総合研究大学院大学、6. 日本流星研究会、7. 国立天文台、8. 東京大学大学院理学系研究科 天文学教室、9. 東京大学ビッグバン宇宙国際研究センター、10. 東北大学 天文学教室、11. 京都大学、12. 甲南大学、13. 統計数理研究所、14. 日本スペースガード協会

1. Institute of Astronomy, Graduate School of Science, The University of Tokyo, 2. Koyama Astronomical Observatory, Kyoto Sangyo University, 3. Center for Planetary Science, c/o Integrated Research Center of Kobe University, 4. Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 5. The Graduate University for Advanced Studies (SOKENDAI), 6. The Nippon Meteor Society, 7. National Astronomical Observatory of Japan, 8. Department of Astronomy, Graduate School of Science, The University of Tokyo, 9. Research Center for the Early Universe, the University of Tokyo, 10. Astronomical Institute, Tohoku University, 11. Kyoto University, 12. Konan University, 13. The Institute of Statistical Mathematics, 14. Japan Spaceguard Association

There are vastly different sizes of objects in the solar system, from the Sun, which is obviously the largest object in the solar system, to beta-meteoroids, which are small particles pushed away by radiation into interstellar space. The size distribution of small bodies in the solar system is important to understand the origin and the evolution of the solar system. Particles smaller than about 1 mm are usually referred as to interstellar dust particles (IDPs). The IDPs are rapidly (up to a few thousand years) removed by a radiation pressure or the Poynting-Robertson effect. Present activities of solar system small bodies are reflected in the size distribution of the IDPs. The IDPs in interplanetary space are, however, difficult to investigate directly, since they are too small to be observed and too sparsely distributed for in-situ investigation.

Tons of IDPs fall down to the Earth per day. Interacting with atmosphere, they lose part of their kinetic energy in a form of emission. This phenomenon is called meteor. Thus, we can derive physical parameters of single particles by investigating meteors. The size distribution of the IDPs around the Earth can be examined through the luminosity function of meteors. Fainter meteors should be observed to constrain the size distribution of smaller IDPs.

To efficiently detect faint meteors, a high sensitivity video camera with a large aperture is preferred. We have been developing a new wide-field CMOS mosaic camera, Tomo-e Gozen, for the 1.05-m Kiso Schmidt Telescope. The camera is composed of four individual camera modules, each of which is

equipped with 21 CMOS image sensors. Tomo-e Gozen can monitor a sky of about 20 deg^2 continuously at up to 2 Hz. The camera will start its operation with a single module in February, 2018. A limiting magnitude of Tomo-e Gozen is expected to be about 18.5 mag. in the V-band for 0.5 s integration, corresponding to a meteor limiting magnitude of about 13 mag. in the V-band. Tomo-e Gozen mounted on the 1.05-m Kiso Schmidt Telescope will be the world-largest video camera, and thus will be an ideal facility to observe faint meteors. As a pathfinder project for Tomo-e Gozen, we have developed a prototype mosaic CMOS camera, Tomo-e PM. Tomo-e PM is equipped with 8 CMOS sensors, which has an ability to continuously obtain images of about 2 deg^2 at 2 Hz. A limiting magnitude of the Tomo-e PM is as good as of Tomo-e Gozen. Tomo-e PM is still a powerful facility to observe faint meteors.

We performed meteor observations with the Tomo-e PM mounted on the 1.05-m Kiso Schmidt Telescope. The observations were carried out on 2016-04-11 and 2016-04-14. The lunar ages were 3.6 and 6.6 on 2016-04-11 and 2016-04-14, respectively. Net observing time was about 10.6 hours in total. Tomo-e PM detected 2,220 unique meteor events. The video rate absolute magnitudes of the detected meteors ranged from 4.0 to 10.0 mag. in the V-band. The result clearly demonstrates that the Tomo-e Gozen has an ability to investigate meteors fainter than 10 mag. Since no significant meteor shower activity was reported, most of the detected meteors were expected to be sporadic meteors.

We approximate the luminosity function of meteors by an exponential distribution: $\log_{10} N(<M) = \log_{10} N_0 + M \log_{10} r$, where $N(<M)$ is the number of meteors brighter than M -th magnitude, N_0 is the number of meteors brighter than zeroth magnitude, and r is the slope of the luminosity function or the meteor index. The parameters N_0 and r were constrained by a fitting based on a statistical model. The present results suggest that $r = 3.1 \pm 0.4$ and $\log_{10} N_0 = -5.5 \pm 0.5$. The present results are roughly consistent with the luminosity functions obtained for sporadic meteors in literature. We have demonstrated that a wide-field CMOS camera mounted on a large telescope has advantages in observing faint meteors. With Tomo-e Gozen, we are able to efficiently investigate variations in the luminosity function of faint meteors with a sufficiently large number of samples.

キーワード：流星、動画観測、惑星間空間ダスト

Keywords: meteor, video observation, interplanetary dust particle