

地球の鉄コンクリーションの形成メカニズムから読み解く火星の表層環境史

Formation mechanism of Fe-oxide concretions on Earth and its implication for alteration history in early Mars

*長谷川 精¹、吉田 英一²、勝田 長貴³、城野 信一⁴、丸山 一平⁴、南 雅代⁵、浅原 良浩⁴、西本 昌司⁶、山口 靖⁴、Ichinnorov Niiden⁷、Metcalf Richard⁸

*Hitoshi Hasegawa¹, Hidekazu Yoshida², Nagayoshi Katsuta³, Sin-iti Sirono⁴, Ippei Maruyama⁴, Masayo Minami⁵, Yoshihiro Asahara⁴, Shoji Nishimoto⁶, Yasushi Yamaguchi⁴, Niiden Ichinnorov⁷, Richard Metcalfe⁸

1. 高知大学理工学部、2. 名古屋大学博物館、3. 岐阜大学教育学部、4. 名古屋大学環境学研究科、5. 名古屋大学宇宙地球環境研究所、6. 名古屋市科学館、7. モンゴル古生物地質研究所、8. Quintessa Limited, The Hub, Henley-on-Thames, Oxfordshire

1. Faculty of Science and Technology, Kochi University, 2. Nagoya University Museum, 3. Department of Education, Gifu University, 4. Graduate School of Environmental Studies, Nagoya University, 5. Institute of Space-Earth Environmental Research, Nagoya University, 6. Nagoya City Science Museum, 7. Institute of Paleontology and Geology, Mongolian Academy of Science, 8. Quintessa Limited, The Hub, Henley-on-Thames, Oxfordshire

Spherical Fe-oxide concretions on Earth, in particular in Utah, U.S.A, have been investigated as an analogue of hematite spherules discovered in Meridiani Planum on Mars, in order to support interpretations of water-rock interactions in early Mars. Although several formation mechanisms have been proposed for the concretions on Earth and Mars, it is still unclear whether these mechanisms are viable because a precise formation process and precursor of the Fe-oxide concretions are missing. Here, we show evidence that Fe-oxide concretions in Utah and newly discovered Fe-oxide concretions in Mongolia, had spherical calcite (CaCO_3) concretions as precursors. Observed different formation stages of calcite and Fe-oxide concretions, both in the Navajo Sandstone, Utah, and the Djadokhta Formation, Mongolia, indicate the formation process of Fe-oxide concretions as follows: (1) calcite concretions initially formed by groundwater evaporation within aeolian sandstone strata; (2) the calcite concretions were dissolved by infiltrating Fe-rich acidic waters; and (3) mobilized Fe in acidic waters was fixed to form spherical $\text{FeO}(\text{OH})$ (goethite) crusts on the pre-existing spherical calcite concretion surfaces due to the pH-buffering dissolution reaction. The similarity between these Fe-oxide concretions on Earth and the hematite spherule occurrences in Meridiani Planum, combined with evidence of acid sulfate water influences on Mars, suggests that the Martian spherules also formed from dissolution of pre-existing carbonate concretions. Formation of recently discovered spherical-shaped nodules in Gale crater on Mars can also be explained by a similar process, although evidence of acid water influence is not obvious in lower strata of the Gale crater. The hematite spherules in Meridiani Planum and spherical nodules in Gale crater are possibly relics of carbonate minerals formed under a dense thick carbon dioxide atmosphere in the past.

キーワード：コンクリーション、地球、火星、変質、炭酸塩

Keywords: Concretion, Earth, Mars, Alteration, Carbonate