

## Formation of Uranian system via a giant impact

\*上田 翔士<sup>1</sup>、井田 茂<sup>2</sup>、荻原 正博<sup>3</sup>、Morbidei Alessandro<sup>4</sup>、Michel Patrick<sup>4</sup>、Guillot Tristan<sup>4</sup>、細野 七月<sup>5</sup>、斎藤 貴之<sup>2</sup>

\*Shoji Ueta<sup>1</sup>, Shigeru Ida<sup>2</sup>, Masahiro Ogihara<sup>3</sup>, Alessandro Morbidelli<sup>4</sup>, Patrick Michel<sup>4</sup>, Tristan Guillot<sup>4</sup>, Natsuki Hosono<sup>5</sup>, Takayuki R Saitoh<sup>2</sup>

1. 京都大学、2. 東京工業大学、地球生命研究所、3. 国立天文台、4. ニース天文台、5. JAMSTEC

1. Kyoto University, 2. Tokyo Tech, ELSI, 3. NAOJ, 4. Nice Observatory, 5. JAMSTEC

Uranus has 5 regular satellites. The obliquity of Uranus is over 90°, which is anomalous in our solar system. A giant impact scenario would explain origins of these properties. In this study, we perform high-resolution SPH simulations ( $N \sim 10^7$ ) for the formation of Uranian system via Giant Impact and show how much material is released around Uranus. In some cases, we find that solid materials are distributed in circum-Uranian orbits after a giant impact, which is massive enough to create satellites. We also find that the amount of materials that are distributed to the circum-Uranian orbit depends on the equation of state (EoS). We want to show results of high-resolution simulations using EoSs, SESAME and ANEOS.

キーワード：天王星、巨大衝突

Keywords: Uranus, Giant Impact