

## Magmatism at the Outer Rise and Impact of Subduction: SCORE/APL proposals for outer rise drilling

\*山口 飛鳥<sup>1</sup>、藤江 剛<sup>2</sup>、平野 直人<sup>3</sup>、鹿児島 涉悟<sup>1</sup>、秋澤 紀克<sup>1</sup>、森下 知晃<sup>4</sup>、小野 重明<sup>5</sup>、土岐 知弘<sup>6</sup>、山野 誠<sup>7</sup>、木村 純一<sup>8</sup>

\*Asuka Yamaguchi<sup>1</sup>, Gou Fujie<sup>2</sup>, Naoto Hirano<sup>3</sup>, Takanori Kagoshima<sup>1</sup>, Norikatsu Akizawa<sup>1</sup>, Tomoaki Morishita<sup>4</sup>, Shigeaki Ono<sup>5</sup>, Tomohiro Toki<sup>6</sup>, Makoto Yamano<sup>7</sup>, Jun-Ichi Kimura<sup>8</sup>

1. 東京大学大気海洋研究所、2. 海洋研究開発機構地震津波海域観測研究開発センター、3. 東北大学東北アジア研究センター、4. 金沢大学理工研究域自然システム学系、5. 海洋研究開発機構海洋掘削科学研究開発センター、6. 琉球大学理学部海洋自然科学科、7. 東京大学地震研究所、8. 海洋研究開発機構地球内部物質循環研究分野

1. Atmosphere and Ocean Research Institute, The University of Tokyo, 2. Research and Development (R&D) Center for Earthquake and Tsunami (CEAT), Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 3. Center for Northeast Asian Studies, Tohoku University, 4. Faculty of Natural System, Institute of Science and Engineering, Kanazawa University, 5. Research and Development Center for Ocean Drilling Science (ODS), Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 6. Department of Chemistry, Biology, and Marine Science, Faculty of Science, University of the Ryukyus, 7. Earthquake Research Institute, the University of Tokyo, 8. Department of Solid Earth Geochemistry, Japan Agency for Marine-Earth Science and Technology (JAMSTEC)

The Pacific Plate in the NW Pacific includes a few hundred meters-thick pelagic sediment, representing a typical structure of an old oceanic plate. Recent seismic reflection studies, however, showed remarkable variation in the thickness of the sediment layer on the outer rise. The typical thickness of the sediment layer is roughly 300–400 m, whereas considerably wide areas have a sediment thickness of only 50 m or less while their surface topography is not clearly identified. Sedimentation rate of surface sediment in the region is ~5–25 cm/kyr, suggesting that all the sediments have been deposited within recent several million years. We hypothesize that the thin sediment cover is owing to basalt sill intrusions or sheet lava extrusions in and on previous thick pelagic sediment package, and that the intrusive and extrusive basalts might have been fed by petit-spot volcanism occurred around the outer rise. Considering the wide distribution of the acoustic thin sediment cover, we could have vastly underestimated the volume of petit-spot magmatism. If this is the case, the petit-spot magmatism may influence subduction zone seismicity, and therefore global geochemical cycle.

To test this hypothesis, we submitted drilling proposals of J-DESC SCORE (Chikyu Shallow Core Program) and IODP APL (Ancillary Project Letter), both of which we targeted at the outer rise of the Japan Trench. In this presentation, we present background and recent findings related in the two proposals.